

- Glavind L, Lund B, Løe H. The relationship between periodontal state and diabetes duration, insulin dosage and retinal changes. *J Periodontol* 1968 Nov;39(6):341-7.
- Glick M, Garfunkel AA. Common oral findings in two different diseases—leukemia and AIDS. Part 1. *Compendium* 1992 Jun;13(6):432,434,436 passim.
- Glick M, Muzyka BC, Lurie D, Salkin LM. Oral manifestations associated with HIV-related disease as markers for immune suppression and AIDS. *Oral Surg Oral Med Oral Pathol* 1994a Apr;77(4):344-9.
- Glick M, Muzyka BC, Salkin LM, Lurie D. Necrotizing ulcerative periodontitis: a marker for immune deterioration and a predictor for the diagnosis of AIDS. *J Periodontol* 1994b May;65(5):393-7.
- Gonzalez M, Banderas JA, Baez A, Belmont R. Salivary lead and cadmium in a young population residing in Mexico City. *Toxicol Lett* 1997 Sep 19;93(1):55-64.
- Goteiner D, Sonis ST, Fasciano R. Cavernous sinus thrombosis and brain abscess initiated and maintained by periodontally involved teeth. *J Oral Med* 1982;37:80-3.
- Goteiner D, Vogel R, Deasy M, Goteiner C. Periodontal and caries experience in children with insulin-dependent diabetes mellitus. *J Am Dent Assoc* 1986 Aug;113(2):277-9.
- Grau AJ, Buggle F, Ziegler C, Schwarz W, Meuser J, Tasman AJ, Buhler A, Benesch C, Becher H, Hacke W. Association between acute cerebrovascular ischemia and chronic and recurrent infection. *Stroke* 1997 Sep;28(9):1724-9.
- Greenberg MS. HIV-associated lesions. *Dermatol Clin* 1996 Apr;14(2):319-26.
- Greenspan D, Greenspan JS. HIV-related oral disease. *Lancet* 1996 Sep;348(9029):729-33.
- Greenspan D, Greenspan JS, Conant M, Petersen V, Silverman S Jr, De Souza Y. Oral "hairy" leukoplakia in male homosexuals: evidence of association with both papillomavirus and a herpes-group virus. *Lancet* 1984 Oct 13;2(8407):831-4.
- Greenspan D, Greenspan JS, Hearst NG, Pan LZ, Conant MA, Abrams DI, Hollander H, Levy JA. Relation of oral hairy leukoplakia to infection with the human immunodeficiency virus and the risk of developing AIDS. *J Infect Dis* 1987 Mar;155(3):475-81.
- Greenspan D, Greenspan JS, Souza YG, Levy JA, Ungar AM. Oral hairy leukoplakia in an HIV-negative renal transplant recipient. *J Oral Pathol Med* 1989 Jan;18(1):32-4.
- Greenspan JS. In: Fauci AS, Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Kasper DI, editors. *Harrison's principles of internal medicine*. 14th ed. New York: McGraw-Hill; 1998.
- Greenspan JS, Greenspan D, Lennette ET, Abrams DI, Conant MA, Petersen VH. Oral viral leukoplakia—a new AIDS-associated condition. *Adv Exp Med Biol* 1985;187:123-8.
- Grossi SG, Genco RJ. Periodontal disease and diabetes mellitus: a two-way relationship. *Ann Periodontol* 1998 Jul;3(1):51-61.
- Grossi SG, Zambon JJ, Ho AW, Koch G, Dunford RG, Machtei EE, Norderyd OM, Genco RJ. Assessment of risk for periodontal disease. I. Risk indicators for attachment loss. *J Periodontol* 1994 Mar;65(3):260-7.
- Grossi SG, Skrepcinski FB, DeCaro T, Zambon JJ, Cummins D, Genco RJ. Response to periodontal therapy in diabetics and smokers. *J Periodontol* 1996 Oct;67(10 Suppl):1094-102.
- Grossi SG, Skrepcinski FB, DeCaro T, Robertson DC, Ho AW, Dunford RG, Genco RJ. Treatment of periodontal disease in diabetes reduces glycated hemoglobin. *J Periodontol* 1997 Aug;68(8):713-9.
- Grossi SG, Donahue R, Quattrin T, Trevisan M, Ho A, Genco RJ. Periodontal disease increases the risk for insulin resistance. *J Dent Res* 1999;78(IADR abstract no. 3486).
- Guyen Y, Satman I, Dinccag N, Alptekin S. Salivary peroxidase activity in whole saliva of patients with insulin-dependent (type-1) diabetes mellitus. *J Clin Periodontol* 1996 Sep;23(9):879-81.
- Haeckel R. Factors influencing the saliva/plasma ratio of drugs. In: Malamud D, Tabak L, editors. *Saliva as a diagnostic fluid*. Ann NY Acad Sci 1993;694:128-42.
- Hallgren R, Lundquist G. Elevated serum levels of pancreatic polypeptide are related to impaired glucose handling in inflammatory states. *Scand J Gastroenterol* 1983 May;18(4):561-4.
- Harrison R, Bowen WH. Periodontal health, dental caries, and metabolic control in insulin-dependent diabetic children and adolescents. *Pediatr Dent* 1987 Dec;9(4):283-6.
- Hayes C, Sparrow D, Cohen M, Vokonas PS, Garcia RI. The association between alveolar bone loss and pulmonary function: the VA Dental Longitudinal Study. *Ann Periodontol* 1998 Jul;3(1):257-61.
- Heimdahl A, Gahrton G, Groth G-C, et al. Selective decontamination of the alimentary tract microbial flora in patients treated with bone marrow transplantation. *Scand J Infect Dis* 1984;16(1):51-60.
- Heimdahl A, Mattson T, Dahllöf F, Lonnquist B, Ringden O. The oral cavity as a port of entry for early infections in patients treated with bone marrow transplantation. *Oral Surg Oral Med Pathol* 1989;68:711-6.
- Herzberg MC, Meyer MW. Effects of oral flora on platelets: possible consequences in cardiovascular disease. *J Periodontol* 1996 Oct;67(10 Suppl):1138-42.
- Herzberg MC, Meyer MW. Dental plaque, platelets, and cardiovascular diseases. *Ann Periodontol* 1998 Jul;3(1):151-60.
- Herzberg MC, Brintzenhofe KL, Clawson CC. Aggregation of human platelets and adhesion of *Streptococcus sanguis*. *Infect Immun* 1983 Mar;39(3):1457-69.

- Herzberg MC, MacFarlane GD, Liu P, Erickson PR. The platelet as an inflammatory cell in periodontal diseases: interactions with *Porphyromonas gingivalis*. In: Genco R, Hamada S, Lehner T, McGhee J, Mergenhagen S, editors. Molecular pathogenesis of periodontal disease. Washington: American Society for Microbiology; 1994. p. 247-55.
- Hildebolt CF, Pilgram TK, Dotson M, Yokoyama-Crothers N, Muckerman J, Hauser J, Cohen S, Kardaris E, Vannier MW, Hanes P, Shrout MK, Civitelli R. Attachment loss with postmenopausal age and smoking. *J Periodontol* 1997 Oct;32(7):619-25.
- Hillier SL, Martius J, Krohn MJ, Kiviat N, Holmes KK, Eschenbach DA. A case-control study of chorioamniotic infection and histologic chorioamnionitis in prematurity. *N Engl J Med* 1988 Oct;319(15):972-8.
- Hillier SL, Nugent RP, Eschenbach DA, Krohn MA, Gibbs RS, Martin DH, Cotch MF, Edelman R, Pastorek JG 2nd, Rao AV, et al. Association between bacterial vaginosis and preterm delivery of a low-birth-weight infant. The vaginal infections and prematurity study group. *N Engl J Med* 1995 Dec;333(26):1737-42.
- Hirai T, Ishijima T, Hashikawa Y, Yajima T. Osteoporosis and reduction of residual ridge in edentulous patients. *J Prosthet Dent* 1993 Jan;69(1):49-56.
- Holmes K, Sparling PF, Mardh P, Lemon SM, Stamm W, Piot P, Wasserheit J, editors. Sexually transmitted diseases. 3rd ed. New York: McGraw-Hill; 1999.
- Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science* 1993 Jan;259(5091):87-91.
- Hove KA, Stallard RE. Diabetes and the periodontal patient. *J Periodontol* 1970 Dec;41(12):713-8.
- Hugoson A, Thorstensson H, Falk H, Kuylenstierna J. Periodontal conditions in insulin-dependent diabetics. *J Clin Periodontol* 1989 Apr;16(4):215-23.
- Humphries S, Devlin H, Worthington H. A radiographic investigation into bone resorption of mandibular alveolar bone in elderly edentulous adults. *J Dent* 1989 Apr;17(2):94-6.
- Institute of Medicine (IOM), Committee to Study the Prevention of Low Birth Weight, Division of Health. Promotion and disease progression. Preventing low birth weight. Washington: National Academy Press; 1985.
- Itin P, Ruffli T, Rudlinger R, Cathomas G, Huser B, Podvinec M, Gudat F. Oral hairy leukoplakia in a HIV-negative renal transplant patient: a marker for immunosuppression? *Dermatologica* 1988;177(2):126-8.
- Itin PH, Lautenschlager S, Fluckiger R, et al. Oral manifestations in HIV-infected patients: diagnosis and management. *J Am Acad Dermatol* 1993;29:749-60.
- Jeffcoat MK. Osteoporosis: a possible modifying factor in oral bone loss. *Ann Periodontol* 1998;3:312-21.
- Jeffcoat MK, Lewis CE, Reddy MS, Wang CY, Reford M. Post-menopausal bone loss and its relationships to oral bone loss. *Periodontology* 2000. (in press).
- Joselow MM, Ruiz R, Goldwater LJ. Absorption and excretion of mercury in man. XIV. Salivary excretion of mercury and its relationship to blood and urine mercury. *Arch Environ Health* 1968 Jul;17(1):39-43.
- Joshi N, O'Bryan T, Appelbaum PC. Pleuropulmonary infections caused by *Eikenella corrodens*. *Rev Infect Dis* 1991 Nov-Dec;13(6):1207-12.
- Joshi KJ, Rimm EB, Douglass CW, Trichopoulos D, Asherio A, Willett WC. Poor oral health and coronary heart disease. *J Dent Res* 1996 Sep;75(9):1631-6.
- Joshi KJ, Ascherio A, Rimm E, Douglass CW, Willett WC. The relation between tooth loss and incidence of ischemic stroke. *Circulation* 1999;99:1121.
- Kaye D. Infective endocarditis. In: Fauci AS, Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Kasper DL, editors. *Harrison's principles of internal medicine*. 13th ed. New York: McGraw-Hill; 1994. p. 520-6.
- King GN, Healy CM, Glover MT, Kwan JT, Williams DM, Leigh IM, Thornhill MH. Prevalence and risk factors associated with leukoplakia, hairy leukoplakia, erythematous candidiasis, and gingival hyperplasia in renal transplant recipients. *Oral Surg Oral Med Oral Pathol* 1994 Dec;78(6):718-26.
- Kjellman O, Henriksson CO, Berghagen N, Andersson B. Oral conditions in 105 subjects with insulin-treated diabetes mellitus. *Sven Tandlak Tidsskr* 1970 Feb;63(2):99-110.
- Klein A, Bruser B, Bast M, Rachlis A. Progress of HIV infection and changes in the lipid membrane structure of CD4+ cells. *AIDS* 1992 Mar;6(3):332-3.
- Klein RS, Harris CA, Small CB, Moll B, Lesser M, Friedland GH. Oral candidiasis in high-risk patients as the initial manifestation of the acquired immunodeficiency syndrome. *N Engl J Med* 1984 Aug 9;311(6):354-8.
- Kline MW. Oral manifestations of pediatric human immunodeficiency virus infection: a review of the literature. *Pediatrics* 1996 Mar;97(3):380-8.
- Koelle DM, Huang ML, Chandran B, Vieira J, Piepkorn M, Corey L. Frequent detection of Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) DNA in saliva of human immunodeficiency virus-infected men: clinical and immunologic correlates. *J Infect Dis* 1997;176(1):94-102.
- Krall EA, Dawson-Hughes B, Papas A, Garcia RI. Tooth loss and skeletal bone density in healthy postmenopausal women. *Osteoporos Int* 1994 Mar;4(2):104-9.
- Krall EA, Garcia RI, Dawson-Hughes B. Increased risk of tooth loss is related to bone loss at the whole body, hip, and spine. *Calcif Tissue Int* 1996 Dec;59(6):433-7.

- Kribbs PJ. Comparison of mandibular bone in normal and osteoporotic women. *J Prosthet Dent* 1990 Feb; 63(2):218-22.
- Kribbs PJ, Chesnut CH 3rd. Osteoporosis and dental osteopenia in the elderly. *Gerodontology* 1984 Summer;3(2):101-6.
- Kribbs PJ, Smith DE, Chesnut CH 3rd. Oral findings in osteoporosis. Part II: Relationship between residual ridge and alveolar bone resorption in generalized skeletal osteopenia. *J Prosthet Dent* 1983 Nov;50(5): 719-24.
- Kribbs PJ, Chesnut CH 3rd, Ott SM, Kilcoyne RF. Relationships between mandibular and skeletal bone in an osteoporotic population. *J Prosthet Dent* 1989 Dec;62(6):703-7.
- Kribbs PJ, Chesnut CH 3rd, Ott SM, Kilcoyne RF. Relationships between mandibular and skeletal bone in a population of normal women. *J Prosthet Dent* 1990 Jan;63(1):86-9.
- Kroes I, Lepp PW, Relman DA. Bacterial diversity within the human subgingival crevice. *Proc Natl Acad Sci USA* 1999 Dec 7;96(25):14547-52.
- Lacassin F, Hoen B, Leport C, Selton-Suty C, Delahaye F, Goulet V, Etienne J, Briancon S. Procedures associated with infective endocarditis in adults. A case control study. *Eur Heart J* 1995 Dec;16(12):1968-74.
- Lafferty WE, Hughes JP, Handsfield HH. Sexually transmitted diseases in men who have sex with men. Acquisition of gonorrhea and nongonococcal urethritis by fellatio and implications for STD/HIV prevention. *Sex Transm Dis* 1997 May;24(5):272-8.
- Leggott PL. Oral manifestations in pediatric HIV infection. In: Greenspan JS, Greenspan D, editors. *Oral manifestations of HIV infection*. Chicago: Quintessence; 1995. p. 234-9.
- Lifson AR, Hilton JF, Westenhouse JL, Canchola AJ, Samuel MC, Katz MH, Buchbinder SP, Hessel NA, Osmond DH, Shiboski S, et al. Time from HIV seroconversion to oral candidiasis or hairy leukoplakia among homosexual and bisexual men enrolled in three prospective cohorts. *AIDS* 1994 Jan;8(1):73-9.
- Löe H. Periodontal disease—the sixth complication of diabetes mellitus. *Diabetes Care* 1993 Jan;16(1): 329-34.
- Lorenz KA, Weiss PJ. Capnocytophagal pneumonia in a healthy man. *West J Med* 1994 Jan;160(1):79-80.
- Lucas RM, Howell LP, Wall BA. Nifedipine-induced gingival hyperplasia. A histochemical and ultrastructural study. *J Periodontol* 1985 Apr;56(4):211-5.
- Lucas VS, Roberts GJ, Beighton D. Oral health of children undergoing allogeneic bone marrow transplantation. *Bone Marrow Transpl* 1998;22:801-8.
- Mackenzie RS, Millard HD. Interrelated effects of diabetes, arteriosclerosis and calculus on alveolar bone loss. *J Am Dent Assoc* 1963;66:191-8.
- Malamud D. Oral diagnostic testing for detecting human immunodeficiency virus-1 antibodies: a technology whose time has come. *Am J Med* 1997; 102(4A):9-14.
- Malamud D, Tabak L, editors. *Saliva as a diagnostic fluid*. *Ann NY Acad Sci* 1993;694:128-42.
- Mandel ID. The diagnostic uses of saliva. *J Oral Pathol Med* 1990;19:119-25.
- Manouchehr-Pour M, Bissada NF. Periodontal disease in juvenile and adult diabetic patients: a review of the literature. *J Am Dent Assoc* 1983 Nov;107(5): 766-70.
- Martino R, Subira M, Manteiga R, Badell I, Argiles B, Sureda A, Brunet S. Viridans streptococcal bacteremia and viridans streptococcal shock syndrome in neutropenic patients: comparison between children and adults receiving chemotherapy or undergoing bone marrow transplantation. *Clin Infect Dis* 1995;20:476-7.
- Mattila KJ, Nieminen MS, Valtonen VV, Rasi VP, Kesaniemi YA, Syrjala SL, Jungell PS, Isoluoma M, Hietaniemi K, Jokinen MJ. Association between dental health and acute myocardial infarction. *BMJ* 1989 Mar 25;298(6676):779-81.
- Mattila KJ, Valtonen VV, Nieminen M, Huttunen JK. Dental infection and the risk of new coronary events: prospective study of patients with documented coronary artery disease. *Clin Infect Dis* 1995 Mar;20(3): 588-92.
- Mattila KJ, Asikainen S, Wolf J, Jousimies-Somer H, Valtonen VV, Nieminen M. Age, dental infections, and coronary heart disease. *J Dent Res* 2000;79: 756-60.
- McCormick MC. The contribution of low birth weight to infant mortality and childhood morbidity. *N Engl J Med* 1985 Jan;312(2):82-90.
- Mealey BL. Periodontal implications: medically compromised patients. *Ann Periodontol* 1996;1:256-321.
- Meyer DH, Sreenivasan PK, Fives-Taylor PM. Evidence for invasion of a human oral cell line by *Actinobacillus actinomycetemcomitans*. *Infect Immun* 1991; 59:2719-26.
- Miller LS, Manwell MA, Newbold D, Reding ME, Rasheed A, Blodgett J, Kornman KS. The relationship between reduction in periodontal inflammation and diabetes control: a report of 9 cases. *J Periodontol* 1992 Oct;63(10):843-8.
- Mohajery M, Brooks SL. Oral radiographs in the detection of early signs of osteoporosis. *Oral Surg Oral Med Oral Pathol* 1992 Jan;73(1):112-7.
- Montaner JS, Le TN, Le N, Craib KJ, Schechter MT. Application of the World Health Organization system for HIV infection in a cohort of homosexual men in developing a prognostically meaningful staging system. *AIDS* 1992 Jul;6(7):719-24.
- Morfeldt-Manson L, Julander I, Nilsson B. Dermatitis of the face, yellow toe nail changes, hairy leukoplakia and oral candidiasis are clinical indicators of progression to AIDS/opportunistic infection in patients with HIV infection. *Scand J Infect Dis* 1989;21(5):497-505.
- Morris JF, Sewell DL. Necrotizing pneumonia caused by mixed infection with *Actinobacillus actinomycetem-*

- comitans* and *Actinomyces israelii*: case report and review. *Clin Infect Dis* 1994 Mar;18(3):450-2.
- Morrison HI, Ellison LF, Taylor GW. Periodontal disease and risk of fatal coronary heart and cerebrovascular diseases. *J Cardiovasc Risk* 1999;6:7-11.
- Morton AA, Williams RW, Watts TL. Initial study of periodontal status in non-insulin-dependent diabetics in Mauritius. *J Dent* 1995 Dec;23(6):343-5.
- Mouldsdale MT, Eykyn SJ, Phillips I. Infective endocarditis, 1970-1979. A study of culture-positive cases in St. Thomas' Hospital. *Q J Med* 1980;49(195):315-28.
- Murrah VA. Diabetes mellitus and associated oral manifestations: a review. *J Oral Pathol* 1985 Apr;14(4):271-81.
- Murray CG, Herson J, Daly TE, Zimmerman S. Radiation necrosis of the mandible: a 10-year study. Part I. Factors influencing the onset of necrosis. *Int J Radiat Oncol Biol Phys* 1980a;6:543-8.
- Murray CG, Herson J, Daly TE, Zimmerman S. Radiation necrosis of the mandible: a 10-year study. Part II. Dental factors; onset, duration and management of necrosis. *Int J Radiat Oncol Biol Phys* 1980b;6:549-53.
- National Institute of Arthritis, Musculoskeletal and Skin Diseases. Available from: <http://www.nih.gov/niams/healthinfo/opbkggr.htm> [2000].
- National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Diabetes statistics. Bethesda (MD): National Institutes of Health; 1999 Mar. NIH Pub. no. 99-3892.
- National Institutes of Health (NIH). Consensus Development Conference on Oral Complication of Cancer Therapy, Diagnosis, Prevention and Treatment. NIH monograph no. 9. Bethesda (MD): National Institutes of Health; 1990.
- Nelson RG, Shlossman M, Budding LM, Pettitt DJ, Saad MF, Genco RJ, Knowler WC. Periodontal disease and NIDDM in Pima Indians. *Diabetes Care* 1990 Aug;13(8):836-40.
- Njoroge T, Genco RJ, Sojar HT, Hamada N, Genco CA. A role for fimbriae in *Porphyromonas gingivalis* invasion of oral epithelial cells. *Infect Immun* 1997 May; 65(5):1980-4.
- Novaes Junior AB, Pereira AL, de Moraes N, Novaes AB. Manifestations of insulin-dependent diabetes mellitus in the periodontium of young Brazilian patients. *J Periodontol* 1991 Feb;62(2):116-22.
- Novaes Junior AB, Gutierrez FG, Novaes AB. Periodontal disease progression in type II non-insulin-dependent diabetes mellitus patients (NIDDM). Part I—Probing pocket depth and clinical attachment. *Braz Dent J* 1996;7(2):65-73.
- O'Farrell BJ, Rajan E, Albloushi SS, Courtney MG, Fielding J, Shattock AG. The reliability of saliva as a sample for the detection of hepatitis A immunoglobulins under various sampling conditions. *Clin Diagn Virol* 1997;7(3):153-7.
- Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, Maynor G, McKaig R, Beck J. Periodontal infection as a possible risk factor for preterm low birth weight. *J Periodontol* 1996 Oct;67(10 Suppl):1103-13.
- Offenbacher S, Jared HL, O'Reilly PG, Wells SR, Salvi GE, Lawrence HP, Socransky SS, Beck JD. Potential pathogenic mechanisms of periodontitis associated pregnancy complications. *Ann Periodontol* 1998 Jul;3(1):233-50.
- Oliver RC, Tervonen T. Periodontitis and tooth loss: comparing diabetics with the general population. *J Am Dent Assoc* 1993 Dec;124(12):71-6.
- Oliver RC, Tervonen T. Diabetes—a risk factor for periodontitis in adults? *J Periodontol* 1994 May;65(5 Suppl):530-8.
- Ortman LF, Hausmann E, Dunford RG. Skeletal osteopenia and residual ridge resorption. *J Prosthet Dent* 1989 Mar;61(3):321-5.
- Ostrow DG, DiFrancisco W. Why some gay men choose condoms and others switch to "safer" sex. *Int Conf AIDS* 1996;11:260.
- Padian N, Glass S. Transmission of HIV possibly associated with exposure of mucous membrane to contaminated blood. *MMWR Morb Mortal Wkly Rep* 1997;46:620-3.
- Payne B, Zachs NR, Reinhardt RA, et al. The association between estrogen status and alveolar bone density changes in post menopausal women with a history of periodontitis. *J Periodontol* 1997;68:24-31.
- Peterson DE. Research advances in mucositis. *Curr Opin Oncol* 1999;11:261-6.
- Peterson DE, Minah GE, Overholser CD, et al. Microbiology of acute periodontal infection in myelosuppressed cancer patients. *J Clin Oncol* 1987; 5:1461-8.
- Phelan JA. Oral manifestations of human immunodeficiency virus infection. *Med Clin North Am* 1997 Mar;81(2):511-31.
- Pinson M, Hoffman WH, Garnick JJ, Litaker MS. Periodontal disease and type 1 diabetes mellitus in children and adolescents. *J Clin Periodontol* 1995 Feb;22(2):118-23.
- Pizzo PA, Meyers J, Freifeld AG, Walsh T. Infections in the cancer patient. In: DeVita VT, Hellman S, Rosenberg SA, editors. *Cancer: principles and practice of oncology*. 4th ed. Philadelphia: J.B. Lippincott Co.; 1993. p. 2292-337.
- Pope M, Frankel S, Steinman R, Elmore D, Ho D, Marx P. Cutaneous dendritic cells promote replication of immunodeficiency viruses. *Adv Exp Med Biol* 1997; 417:395-9.
- Ramos-Gomez FJ, Hilton JF, Canchola AJ, Greenspan D, Greenspan JS, Maldonado YA. Risk factors for HIV-related oral-facial soft-tissue manifestations in children. *Pediatr Dent* 1996 Mar-Apr; 18(2):121-6.
- Redfield RR, Wright DC, Tramont EC. The Walter Reed staging classification for HTLV-III/LAV infection. *N Engl J Med* 1986 Jan;314(2):131-2.

- Reilly TG, Poxon V, Sanders DS, Elliot TS, Walt RP. Comparison of serum, salivary, and rapid whole blood diagnostic tests for *Helicobacter pylori* and their validation against endoscopy based tests. *Gut* 1997;40(4):454-8.
- Richards AL, Rerrault JG, Garingal LT, Manaloto CR, Sie A, Graham R, Ramos RM, Leonardo JB, Hyams KC. A non-invasive assessment of hepatitis B virus carrier status using saliva samples. *Southeast Asian J Trop Med Public Health* 1996;27(1):80-4.
- Ringelberg ML, Dixon DO, Francis AO, Plummer RW. Comparison of gingival health and gingival crevicular fluid flow in children with and without diabetes. *J Dent Res* 1977 Feb;56(2):108-11.
- Riviere GR, Weisz KS, Adams DF, Thomas DD. Pathogen-related oral spirochetes from dental plaque are invasive. *Infect Immun* 1991 Oct;59(10):3377-80.
- Rocke LK, Loprinzi CL, Lee JK, et al. A randomized clinical trial of two different durations of oral cryotherapy for prevention of 5-fluorouracil-related stomatitis. *Cancer* 1993; 72:2234-8.
- Rogers MF, White CR, Sanders R. Lack of transmission of HIV from infected children to their household contacts. *Pediatrics* 1990;85:210-4.
- Rolston KVI, Bodey GP. Infections in patients with cancer. In: Holland JF, Frei E, Bast RC, Kufe DW, Morton DL, Weichselbaum RR, editors. *Cancer medicine*. 3rd ed. Philadelphia: Lea & Febiger; 1993. p. 2416-41.
- Romero R, Baumann P, Gomez R, Salafia C, Rittenhouse L, Barberio D, Behnke E, Cotton DB, Mitchell MD. The relationship between spontaneous rupture of membranes, labor, and microbial invasion of the amniotic cavity and amniotic fluid concentrations of prostaglandins and thromboxane B2 in term pregnancy. *Am J Obstet Gynecol* 1993 Jun;168(6 Pt 1):1654-64; discussion 1664-8.
- Rothenberg RR, Scarlett MI, delRio C, Reznik D, Daniels C. Oral transmission of HIV. *AIDS* 1998;12:2095-105.
- Roy R, Burghoff R, Steffens DL. Detection of STR, D1S80, and gender alleles from body fluids collected on isocode (TM) foam collectors and paper-based devices. *Crime Lab Digest* 1997;24(1):4-9.
- Royce RA, Luckmann RS, Fusaro RE, Winkelstein W Jr. The natural history of HIV-1 infection: staging classifications of disease. *AIDS* 1991 Apr;5(4):355-64.
- Ruescher TJ, Sodeifi A, Scrivan SJ, Kaban LB, Sonis ST. The impact on alpha-hemolytic streptococcal infection in patients undergoing autologous bone marrow transplantation for hematologic malignancies. *Cancer* 1998;82:2275-81.
- Ruprecht RM, Baba TW, Liska V, Ray NB, Martin LN, Murphey-Corb M, Rizvi TA, Bernacky BJ, Keeling ME, McClure HM, Andersen J. Oral transmission of primate lentiviruses. *J Infect Dis* 1999 May;179 (Suppl 3):S408-12.
- Rylander H, Ramberg P, Blohme G, Lindhe J. Prevalence of periodontal disease in young diabetics. *J Clin Periodontol* 1987 Jan;14(1):38-43.
- Saah AJ, Munoz A, Kuo V, Fox R, Kaslow RA, Phair JP, Rinaldo CR Jr, Detels R, Polk BF. Predictors of the risk of development of acquired immunodeficiency syndrome within 24 months among gay men seropositive for human immunodeficiency virus type 1: a report from the Multicenter AIDS Cohort Study. *Am J Epidemiol* 1992;135:1147-55.
- Saal CJ, Mason JC, Cheuk SI, Hill MK. Brain abscess from chronic odontogenic cause: report of case. *J Am Dent Assoc* 1988;117:453-5.
- Saglie FR, Marfany A, Camargo P. Intra-gingival occurrence of *Actinobacillus actinomycetemcomitans* and *Bacteroides gingivalis* in active destructive periodontal lesions. *J Periodontol* 1988 Apr;59(4):259-65.
- Salvi GE, Lawrence HP, Offenbacher S, Beck JD. Influence of risk factors on the pathogenesis of periodontitis. *Periodontol* 2000 1997 Jun;14:173-201.
- Sammalkorpi K. Glucose intolerance in acute infections. *J Intern Med* 1989 Jan;225(1):15-9.
- Sande MA. Transmission of AIDS. The case against casual contagion. *N Engl J Med* 1986;314:380-2.
- Sandler HC, Stahl SS. Prevalence of periodontal disease in a hospitalized population. *J Dent Res* 1960 May-Jun;39(3):439-49.
- Sandros J, Papapanou PN, Dahlen G. *Porphyromonas gingivalis* invades oral epithelial cells in vitro. *J Periodont Res* 1993 May;28(3):219-26.
- Sandros J, Papapanou PN, Nannmark U, Dahlen G. *Porphyromonas gingivalis* invades pocket epithelium in vitro. *J Periodont Res* 1994 Jan;29(1):62-9.
- Scannapieco FA. Position paper of the American Academy of Periodontology: periodontal disease as a potential risk factor for systemic diseases. *J Periodontol* 1998 Jul;69(7):841-50.
- Scannapieco FA. Role of oral bacteria in respiratory infection. *J Periodontol* 1999;70:793-802.
- Scannapieco FA, Mylotte JM. Relationships between periodontal disease and bacterial pneumonia. *J Periodontol* 1996 Oct;67(10 Suppl):1114-22.
- Scannapieco FA, Papandonatos GD, Dunford RG. Associations between oral conditions and respiratory disease in a national sample survey population. *Ann Periodontol* 1998 Jul;3(1):251-6.
- Schmidt MI, Duncan BB, Sharrett AR, Lindberg G, Savage PJ, Offenbacher S, Azambuja MI, Tracy RP, Heiss G. Markers of inflammation and prediction of diabetes mellitus in adults (Atherosclerosis Risk in Communities study): a cohort study. *Lancet* 1999 May 15;353:1649-52.
- Schubert MM, Epstein JB, Peterson DE. Oral complications of cancer therapy. In: Yagelia JA, Neidle EA, Dowd FJ, editors. *Pharmacology and therapeutics in dentistry*. St. Louis: Mosby-Year Book Inc.; 1998. p. 644-55.
- Schubert MM, Peterson DE, Lloid M. Oral complications. In: Thomas E, Blume KG, Forman SJ, editors.

- Hematopoietic cell transplantation. 2nd ed. Malden (MA): Blackwell Science, Inc.; 1999. p. 751-63.
- Schwarzc SK, Kellogg TA, Kohn RP, et al. Temporal trends in human immunodeficiency virus seroprevalence and sexual behavior at the San Francisco municipal sexually transmitted disease clinic, 1989-1992. *Am J Epidemiol* 1995;142:314-22.
- Scully C, Epstein JB. Oral health care for the cancer patient. *Eur J Cancer B Oral Oncol* 1996 Sep;32B(5):281-92.
- Seage GR 3rd, Gatsonis C, Weissman JS, Haas JS, Craven DE, Makadon H, Goldberg J, Coltin K, Lewin KS, Epstein AM, et al., The Boston AIDS Survival Score (BASS): a multidimensional AIDS severity instrument. *Am J Public Health* 1997 Apr;87(4):567-73.
- Semenciw R, Morrison H, Mao Y, Johansen H, Davies JW, Wigle D. Major risk factors for cardiovascular disease mortality in adults: results from the Nutrition Canada Survey. *Int J Epidemiol* 1988;17:317-24.
- Seppala B, Ainamo J. A site-by-site follow-up study on the effect of controlled versus poorly controlled insulin-dependent diabetes mellitus. *J Clin Periodontol* 1994 Mar;21(3):161-5.
- Seppala B, Seppala M, Ainamo J. A longitudinal study on insulin-dependent diabetes mellitus and periodontal disease. *J Clin Periodontol* 1993 Mar;20(3):161-5.
- Shapiro S, McCormick MC, Starfield BH, Krischer JP, Bross D. Relevance of correlates of infant deaths for significant morbidity at 1 year of age. *Am J Obstet Gynecol* 1980 Feb 1;136(3):363-73.
- Shinzato T, Saito A. A mechanism of pathogenicity of "Streptococcus milleri group" in pulmonary infection: synergy with an anaerobe. *J Med Microbiol* 1994 Feb;40(2):118-23.
- Shlossman M, Knowler WC, Pettitt DJ, Genco RJ. Type 2 diabetes mellitus and periodontal disease. *J Am Dent Assoc* 1990 Oct;121(4):532-6.
- Silver JG, Martin AW, McBride BC. Experimental transient bacteraemias in human subjects with varying degrees of plaque accumulation and gingival inflammation. *J Clin Periodontol* 1977 May;4(2):92-9.
- Smith GT, Greenbaum CJ, Johnson BD, Persson GR. Short-term responses to periodontal therapy in insulin-dependent diabetic patients. *J Periodontol* 1996 Aug;67(8):794-802.
- Sonis ST. Oral mucositis as a biological process: a new hypothesis for the development of chemotherapy induced stomatotoxicity. *Oral Oncol* 1998;34:39-43.
- Sonis ST, Van Vugt AG, Brien JP, Muska AD, Bruskin AM, Rose A, Haley JD. Transforming growth factor-beta 3 mediated modulation of cell cycling and attenuation of 5-fluorouracil-induced oral mucositis. *Oral Oncol* 1997 Jan;33(1):47-54.
- Sparrelid E, Häggglund H, Remberger M, Ringden O, Lonnquist B, Ljungman P, Andersson J. Bacteraemia during the aplastic phase after allogeneic bone marrow transplantation is associated with early death from invasive fungal infection. *Bone Marrow Transpl* 1998;22:795-800.
- Sreebny LM, Schwartz SS. A reference guide to drugs and dry mouth. 2nd ed. *Gerodontology* 1997;14:33-47.
- Striskandan S, Soto A, Evans TJ, Cohen J. Viridans streptococcal bacteraemia: a clinical study. *QJM* 1995;88:415-20.
- Stahl-Hennig C, Steinman RM, Tenner-Racz K, Pope M, Stolte N, Matz-Rensing K, Grobschupff G, Raschdorff B, Hunsmann G, Racz P. Rapid infection of oral mucosal-associated lymphoid tissue with simian immunodeficiency virus. *Science* 1999 Aug 20;285(5431):1261-5.
- Strom BL, Abrutyn E, Berlin JA, et al. Dental and cardiac risk factors for infective endocarditis. *Arch Intern Med* 1998 Nov 15;129(10):761-9.
- Svenson KL, Lundquist G, Wide L, Hallgren R. Impaired glucose handling in active rheumatoid arthritis: relationship to the secretion of insulin and counter-regulatory hormones. *Metabolism* 1987 Oct;36(10):940-3.
- Sznajder N, Carraro JJ, Rugna S, Sereday M. Periodontal findings in diabetic and nondiabetic patients. *J Periodontol* 1978 Sep;49(9):445-8.
- Szpunar SM, Ismail AI, Eklund SA. Diabetes and periodontal disease: analyses of NHANES I and HHANES. *J Dent Res* 1989;68(SI):164-438 (abstract 1605).
- Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M, Knowler WC, Pettitt DJ. Severe periodontitis and risk for poor glycemic control in patients with non-insulin-dependent diabetes mellitus. *J Periodontol* 1996 Oct;67(10 Suppl):1085-93.
- Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M, Knowler WC, Pettitt DJ. Non-insulin dependent diabetes mellitus and alveolar bone loss progression over 2 years. *J Periodontol* 1998a Jan;69(1):76-83.
- Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M. Glycemic control and alveolar bone loss progression in type 2 diabetes. *Ann Periodontol* 1998b Jul;3(1):30-9.
- Tervonen T, Karjalainen K. Periodontal disease related to diabetic status. A pilot study of the response to periodontal therapy in type 1 diabetes. *J Clin Periodontol* 1997 Jul;24(7):505-10.
- Tervonen T, Knuuttila M. Relation of diabetes control to periodontal pocketing and alveolar bone level. *Oral Surg Oral Med Oral Pathol* 1986 Apr;61(4):346-9.
- Thorstensson H, Hugoson A. Periodontal disease experience in adult long-duration insulin-dependent diabetics. *J Clin Periodontol* 1993 May;20(5):352-8.
- Tindall B, Carr A, Cooper DA. Primary HIV infection: clinical, immunologic, and serologic aspects. In: Sande MA, Volberding PA, editors. *The medical management of AIDS*. Philadelphia: W.B. Saunders; 1995. p. 105-29.

- U.S. Department of Health and Human Services (USDHHS), Center for Research for Mothers and Children. Public Health Service/National Institutes of Health Progress Report. Washington: U.S. Department of Health and Human Services; 1984.
- U.S. Preventive Services Task Force. Guide to clinical preventive services. 2nd ed. Washington: US GPO; 1996.
- van Schie RC, Wilson ME. Saliva: a convenient source of DNA for analysis of bi-allelic polymorphisms of Fcg receptor IIA (CD32) and Fcg receptor IIIB (CD16). *J Immunol Methods* 1997;208:91-101.
- von Wowern N. Dual-photon absorptiometry of mandibles: in vitro test of a new method. *Scand J Dent Res* 1985 Apr;93(2):169-77.
- von Wowern N. Bone mineral content of mandibles: normal reference values—rate of age-related bone loss. *Calcif Tissue Int* 1988 Oct;43(4):193-8.
- von Wowern N, Klausen B, Kollerup G. Osteoporosis: a risk factor in periodontal disease. *J Periodontol* 1994 Dec;65(12):1134-8.
- Wactawski-Wende J, Grossi SG, Trevisan M, Genco RJ, Tezal M, Dunford RG, Ho AW, Hausmann E, Hreshchyshyn MM. The role of osteopenia in oral bone loss and periodontal disease. *J Periodontol* 1996;67(10 Suppl):1076-84.
- Wallace JI, Bloch D, Whitmore R, et al. Fellatio is a significant risk activity for acquiring AIDS in New York City streetwalking sex workers. *Int Conf AIDS* 1996;11:381.
- Ward JW, Duchin JS. The epidemiology of HIV and AIDS in the United States. *AIDS Clin Rev* 1997;45.
- Weinberg A, Belton CM, Park Y, Lamont RJ. Role of *Porphyromonas gingivalis* fimbriae invasion of gingival epithelial cells. *Infect Immun* 1997 Jan;65(1):313-6.
- Weinstein L, Schlesinger JJ. Pathoanatomic, pathophysiologic and clinical correlations in endocarditis (second of two parts). *New Engl J Med* 1974;291:832-6, 1122-6.
- Westfelt E, Rylander H, Blohme G, Jonasson P, Lindhe J. The effect of periodontal therapy in diabetics. Results after 5 years. *J Clin Periodontol* 1996 Feb;23(2):92-100.
- Wilkes JD. Prevention and treatment of oral mucositis following cancer chemotherapy. *Semin Oncol* 1998;25:538-51.
- Williams ME, Quesenberry PJ. Hematopoietic growth factors. *Hematol Pathol* 1992;6(3):105-24.
- Williams RC Jr, Mahan CJ. Periodontal disease and diabetes in young adults. *JAMA* 1960 Feb 20;172:776-8.
- Wilson JT. Clinical correlates of drugs in saliva. *Ann NY Acad Sci* 1993 Sep 30;694:48-61.
- Wilton JM, Griffiths GS, Curtis MA, Maiden MFJ, Gillet IR, Wilson DT, Sterne JA, Johnson NW. Detection of high-risk groups and individuals for periodontal diseases. Systemic predisposition and markers of general health. *J Clin Periodontol* 1988 Jul;15(6):339-46.
- Winkler JR, Robertson PB. Periodontal disease associated with HIV infection. *Oral Surg Oral Med Oral Pathol* 1992 Feb;73(2):145-50.
- Winkler JR, Grassi M, Murray PA. Clinical description and etiology of HIV-associated periodontal diseases. In: Robertson PB, Greenspan JS, editors. Perspectives on oral manifestations of AIDS: diagnosis and management of HIV-associated infections. Littleton (MA): PSG Publishing; 1988.
- Wolf J. Dental and periodontal conditions in diabetes mellitus. A clinical and radiographic study. *Proc Finn Dent Soc* 1977;73 (4-6 Suppl):1-56.
- Wu T, Trevisan M, Genco RJ, Dorn JP, Falkner KL, Sempos CT. Periodontal disease and risk of cerebrovascular disease: a prospective study of a representative sample of U.S. adults. *Am J Epidemiol* 1999;149:(11)290, Suppl S Jun 1.
- Wu TJ, Trevisan M, Genco RJ, Dorn JP, Falkner KL, Sempos CT. Examination of the relation between periodontal health status and cardiovascular risk factors: serum total and high density lipoprotein, cholesterol, C-reactive protein, and plasma fibrinogen. *Am J Epidemiol* 2000;151:273-82.
- Yavuzylmaz E, Yumak O, Akdoganli T, Yamalik N, Ozer N, Ersoy F, Yenijay I. The alterations of whole saliva constituents in patients with diabetes mellitus. *Aust Dent J* 1996 Jun;41(3):193-7.
- Yuan A, Yang PC, Lee LN, Chang DB, Kuo SH, Luh KT. *Actinobacillus actinomycetemcomitans* pneumonia with chest wall involvement and rib destruction. *Chest* 1992 May;101(5):1450-2.
- Zakrzewska JM, Aly Z, Speight PM. Oral hairy leukoplakia in a HIV-negative asthmatic patient on systemic steroids. *J Oral Pathol Med* 1995 Jul;24(6):282-4.
- Zijlstra EE, Swart GR, Godfroy FJ, Degener JE. Pericarditis, pneumonia and brain abscess due to a combined *Actinomyces-Actinobacillus actinomycetemcomitans* infection. *J Infect* 1992 Jul;25(1):83-7.

Effects on Well-being and Quality of Life

Fifty years ago the World Health Organization (WHO) defined health as the “complete state of physical, mental, and social well-being and not merely the absence of infirmity” (WHO 1948). In its definition the WHO acknowledged that an individual who is technically “cured” of disease may not necessarily be “well” and went on to indicate three dimensions of well-being. *Physical well-being* assumes the ability to function normally in activities such as bathing, dressing, eating, and moving around. *Mental well-being* implies that cognitive faculties are intact and that there is no burden of fear, anxiety, stress, depression, or other negative emotions. *Social well-being* relates to one’s ability to participate in society, fulfilling roles as family member, friend, worker, or citizen or in other ways engaging in interactions with others.

The WHO declaration resonated with ongoing developments in the social sciences as theoreticians recognized the need for multiple indicators in assessing health and treatment outcomes (Bergner et al. 1981, Fries et al. 1982, Hunt et al. 1985, Meenan et al. 1980). These efforts led to definitions of “health-related quality of life” (Guyatt et al. 1993) as well as explanatory models. The model proposed by Wilson and Cleary (1995), for example, posits five dimensions by which to measure treatment outcomes: biological and physiological variables, symptom status, functional status, general health perceptions, and overall quality of life. These factors are not independent but may be reciprocally connected. For example, a diabetic patient with symptoms of depression may experience a rise in serum glucose as a result of less vigilant glucose monitoring; the depression may then lead to a deterioration in physical and social activities. Most importantly, measures of biological and physiological factors are often inconsistent with patients’ own reports of symptoms, ability to function, general health perceptions, and overall

quality of life. In the wake of these developments in general medicine, researchers began to elaborate multidimensional models of “oral-health-related” quality of life.

The efforts to understand these relationships are particularly relevant given the aging of the population. As Gift and Atchison (1995) stated, measuring health-related quality of life allows assessment of “the trade-off between how long and how well people live.” Diseases and disorders that result in dental and craniofacial defects can thwart that goal, disturbing self-image, self-esteem, and well-being. Oral-facial pain and loss of sensorimotor functions limit food choices and the pleasures of eating, restrict social contact, and inhibit intimacy.

Oral complications of many systemic diseases also compromise the quality of life. Problems with speaking, chewing, taste, smell, and swallowing are common in neurodegenerative conditions such as Parkinson’s disease; oral complications of AIDS include pain, dry mouth, mucosal infections, and Kaposi’s sarcoma; cancer therapy can result in painful ulcers, mucositis, and rampant dental caries; and periodontal disease is a complication of diabetes and osteoporosis. Prescription and nonprescription drugs often have the side effect of dry mouth.

The ability to measure the quality of life has the practical value of guiding policymakers, health service researchers, epidemiologists, program evaluators, and clinicians interested in the effects of interventions. The measures can also provide useful information to patients and family members, third-party payers, and employers. For example, measures of the ability to perform activities of daily living may indicate areas where the patient is able and competent, as well as areas where further therapy may be helpful.

This chapter reviews oral-health-related quality of life findings along functional, psychosocial, and economic dimensions, taking into consideration the

influence of cultural and spiritual values. The results of studies in which investigators asked adults how they value their oral health and whether they are satisfied with their oral health care are included. The study of the association between oral health and quality of life is a relatively new but rapidly growing field. A variety of questionnaires have been designed to assess oral-health-related quality of life, and the chapter concludes with a discussion of their use in surveys and analytic studies, and their potential importance in outcome research.

THE CULTURAL CONTEXT

The determination of the health-related quality of life of an individual is implicitly made against a cultural background that includes a set of values, standards, customs, and traditions associated with a particular society.

Decisions about whether to seek care from a dentist, a physician, or other care provider may be influenced by cultural or ethnic perspectives and understanding (Aday and Forthofer 1992, Andersen and Davidson 1997, Davidson and Andersen 1997, Diehnelt et al. 1990, Kiyak 1993, Lee and Kiyak 1992). Different population groups differ in the way they think about health, and in how they define a health problem, determine its seriousness, and decide whether to seek care. In one cultural setting a painful tooth may be enough to motivate care seeking. In another, bleeding, swelling, or fever may be necessary before care is sought. Similarly, decisions about whether to comply with a suggested treatment regimen, whether to engage in self-care, and whether to return for a follow-up appointment are also culturally influenced.

The anthropology and ethnography literature is rich in references to the ways in which different cultures at different times and places have regarded the human body (Hufford 1992, Kleinman 1979). Cultural beliefs regarding the body, health, and disease are often embedded in religious or spiritual traditions, which in turn may govern how diseases and disorders are regarded and treated. A brief description of Western and non-Western perspectives follows.

Cultural Models

In the medical model typical of Western society the body is partitioned into organs and systems, each with identifiable functions. The body is seen as functioning well unless disease disrupts it. Diseases in themselves are understood to be invariable across

cultures. The medical model has traditionally dichotomized body and mind/soul/spirit—science and magic. Such a perspective sees the body as relatively objective and value-free, immune to nonsomatic influences.

That perspective began to change with the pioneering work of Hans Selye in the 1930s on the importance of stress in health and disease (McEwen 1999). Research in the intervening half century has confirmed the reciprocal connections of the nervous, endocrine, and immune systems, not only in relation to stress, but also in terms of the effects of emotions and cognitive processes on health status.

The model that has emerged as a new paradigm in the study of health and disease incorporates biological with psychological and social factors. This biopsychosocial model is defining agendas for research in such fields as behavioral medicine and psychoneuroimmunology. Social and psychological factors are routinely incorporated into health assessments, the better to describe the quality of life. Other societies hold views of the body strikingly different from the medical model. In some cultures, individuals and their care providers conceive of the body as the union of soul and soma. Illness may occur as a result of a “failure in harmony” or “an imbalance of forces.” Schools of medicine in China, India, and other non-Western societies incorporate such principles into their teaching and practice (Hufford 1992).

Combining Perspectives

All Americans hold culturally influenced perspectives on healing and illness (Henderson et al. 1997), some of which come from more traditional beliefs. Some people accept pain as an inevitable part of illness, a necessary evil, or even punishment for past iniquities or shortcomings and may shun pain-relieving drugs (Zborowski 1952). Many pragmatically combine cultural, folk, complementary, and alternative healing practices with participation in conventional care delivery systems. A recent survey indicates that over 50 percent of Americans sought non-traditional therapies for a number of ailments (Eisenberg et al. 1998).

Traditional beliefs are often comforting and satisfying to individuals (Selikowitz 1994). Certainly, Western culture and science have not always improved the quality of people's lives (Harris et al. 1993). Dietary changes to refined foods have been associated with dental caries (Godson and Williams 1996, Navia 1994), obesity, and other deleterious health changes (Selikowitz 1994). The marketing of tobacco products has added to the burden of cancer

and heart and lung problems worldwide. Migrations from traditional community rural life to urban centers have been associated with family disruption and violence, drug abuse, sexually transmitted diseases, and hypertension in developing countries.

On the other hand, Western science may inform some cultural groups that certain traditional child-rearing practices can be detrimental to oral health (Kolasa 1978, Scheper-Hughes 1990). Early childhood caries is a form of tooth decay with complex etiologies. Researchers studying high rates of infant caries among some cultural groups are exploring the extent to which traditional means of soothing crying babies or handling bedtime routines play a role, as well as investigating prenatal nutrition and transmission of infection from caregiver to child (Febres et al. 1997, Kelly and Bruerd 1987, Ripa 1988, Tinanoff and O'Sullivan 1997).

People who hold different cultural perspectives may distance themselves from Western, scientific worldviews (Lee et al. 1993), a behavior that must be addressed in any program of health promotion and disease prevention. Health professionals who understand indigenous or local healing practices and concepts are better able to motivate patients and thereby enable them to integrate elements from various healing systems (Kleinman 1979).

America is undergoing major demographic changes, with the expectation that at some point before 2050 the white population will no longer represent the majority (Henderson et al. 1997). As these changes occur, cultural elements that now reflect minority groups may become more accepted and dominant. However, cultural values are neither static nor omnipotent in shaping people's lives. Furthermore, individuals within a culture manifest their cultural identity in different ways. Therefore, both the direction of these changes and their effects may be hard to predict.

ORAL-HEALTH-RELATED QUALITY OF LIFE DIMENSIONS

Multiple factors act and interact in determining one's quality of life, as Wilson and Cleary (1995) and others have observed. Thus the idea of assessing quality of life along multiple "dimensions" implies a departure from a simple linear scale with excellent quality of life at one end and greatly diminished quality of life at the other. The following sections explore several dimensions, beginning with effects along functional and psychosocial dimensions and concluding with a discussion of economic effects on quality of life.

Functional Dimensions

Investigators have reported on the effects of dental and craniofacial diseases on the ability to eat and enjoy the full range of dietary choices. The impact of less-than-optimal oral health also has been studied in relation to sleep problems, primarily in relation to oral-facial pain.

Eating

Both dental and systemic diseases can profoundly affect appetite and the ability to eat, and hence can compromise overall health and well-being. Because chronic illness and medications increase in aging populations, these effects may be particularly evident among the frail elderly (Ship et al. 1996). Undernutrition was observed in 50 percent of geriatric residents in a U.S. long-term care facility; in many cases, it was linked to eating and swallowing problems (Keller 1993).

Less severe oral disorders have more subtle effects on functions relating to eating, although the high prevalence of those disorders elevates their relative importance among health problems. For example, data from the National Health and Nutrition Examination Survey III indicate that 33.1 percent of people aged 65 and older have no teeth (Marcus et al. 1996). Furthermore, clinical studies indicate that the masticatory efficiency of replacement teeth is at least 30 to 40 percent lower than that of natural teeth (Idowu et al. 1986). Consistent with these findings, surveys of elderly populations in the United States indicate that self-reported chewing problems affect significant proportions of people. For example, in California 1 percent of Medicare enrollees were unable to swallow comfortably, whereas 37 percent of senior center residents reported trouble biting or chewing foods (Table 6.1).

A number of studies have indicated that having missing teeth is linked to a qualitatively poorer diet. For example, in studies of U.S. veterans (Chauncey et al. 1984), Canadians (Brodeur et al. 1993), and Finns (Ranta et al. 1987), people with impaired dentitions preferred soft, easily chewed foods that were lower in fiber and had lower nutrient density than foods eaten by people with intact dentitions. Quality of life clearly suffers when individuals are forced to limit food choices and the foods chosen do not provide optimal nutrition. For example, they would be hard put to comply with the healthful diet recommendation of "five-a-day" helpings of fiber-rich fruits and vegetables. In the elderly, edentulousness and poor oral health may contribute to significant weight loss,

TABLE 6.1
Prevalence of self-reported eating dysfunction in surveys of elderly Americans

Dysfunction Attributed to Oral Condition(s)	Percentage of Population Group Reporting Dysfunction	
	Elderly Persons in California ^a	
	Senior Center Residents	Medicare Enrollees
Had trouble biting or chewing	37	13
Limited the kinds of foods eaten	23	10
Unable to swallow comfortably	10	1
	Elderly Persons in Florida ^b	
Mouth sometimes dry	39	
Noticed an unpleasant taste in mouth	23	
Unable to chew hard things	19	
Experienced change in sense of taste	9	
Difficulty tasting some foods	6	
Noticed change in sense of smell	5	
	Elderly Persons in North Carolina ^c	
	African Americans	Whites
Difficulty chewing any foods	18	6
Felt sense of taste had worsened	13	3
Uncomfortable eating foods	13	6
Had to avoid eating some foods	10	4
Felt digestion had worsened	8	<1
Had to interrupt meals	6	<1

^a n = 1,842 elderly persons (Atchison and Dolan 1990).

^b n = 390 dentate persons aged 65 and older living in retirement communities (Gilbert et al. 1993).

^c n = 440 persons aged 70 and older in five North Carolina counties (Hunt et al. 1995).

which may affect overall health (Blaum et al. 1995, Ritchie et al. 2000, Sullivan et al. 1993).

Clinical research has demonstrated a general reduction in chewing function as the number of missing teeth increases, even when dentures are worn (Carlsson 1984, Feldman et al. 1980, Helkimo et al. 1978). However, clinical studies in Scandinavia have observed relatively good chewing performance when no more than 12 teeth are missing and the remaining 20 teeth are distributed fairly equally, providing good contact between the upper and the lower jaw (Agerberg and Carlsson 1975). This finding has led some researchers to advocate the concept of a "shortened dental arch" as a treatment goal for older adults, that is, retention of at least 20 well-distributed teeth (Kayser 1981). A consistent finding from research into treatment of tooth loss is that removable dentures produce only partial improvement in chewing performance (Garrett et al. 1996, van der Bilt et al. 1994, Witter et al. 1989). Dentures anchored by implants result in significantly better chewing performance than conventional, removable dentures (Geertman et al. 1996).

Self-reported measures of eating ability, satisfaction with eating, and avoidance of foods are widely

used to evaluate the effects of tooth loss and replacement. These measures capture aspects of eating that are not necessarily reflected in the clinical measures of chewing performance with specific foods (Demers et al. 1996). Such studies of self-reported chewing ability confirm several relationships already noted, including the findings that chewing ability declines as the extent of tooth loss increases and that removable dentures do not fully compensate for the reduction (Hildebrandt et al. 1997, Leake 1990, Slade et al. 1996, Wayler et al. 1984). In a study of male veterans with varying levels of tooth loss, Wayler et al. (1982) reported that levels of taste, smell, texture, perceived ease of chewing, and frequency of ingestion were adversely affected only among people with severely compromised dentitions. However, in some studies of prosthodontic treatment, self-reported chewing ability improved following treatment, even

in the absence of significant change in clinical measures of chewing performance (van der Bilt et al. 1994).

Impaired eating due to conditions other than tooth loss has been evaluated less frequently. In a study of dental patients with intact dentitions, Ernest (1993) correlated reduced salivary flow with decreased intake of 18 of 22 nutrients. Locker and Slade (1988) found that 6 percent of the Canadians surveyed who reported symptoms of temporomandibular disorders also reported problems with eating, talking, or swallowing. These conditions are often associated with limited mouth opening and severe pain, which may be constant or present whenever jaw movements are made. The constant dry mouth of patients with Sjögren's syndrome, a disease in which the salivary glands are progressively destroyed, is a major source of discomfort that affects speaking, chewing, and swallowing.

As part of a Performance Status Scale study of oral and pharyngeal cancer patients, List et al. (1990) reported that only one third of patients achieved a perfect score on a measure of normalcy of diet and only 60 percent on a measurement of eating in public. Communication skills also suffered; only 55 percent scored perfectly on understandability of speech.

Sleeping

Sleep problems associated with oral conditions appear to be most closely related to chronic pain, either directly or indirectly in cases where pain and insomnia are exacerbated by depression. In a survey of elderly Floridians, Gilbert et al. (1993) found that 3 percent of the population reported trouble sleeping because of pain or discomfort from dental problems, whereas in a Canadian study Locker and Grushka (1987) reported that 14.2 percent of those with acute or chronic oral-facial pain (or 5.5 percent of the total population) experienced sleep disturbance related to pain. Goulet et al. (1995) assessed sleep problems not specifically attributed to pain among adults in Quebec. The prevalence of sleep problems was 13 percent for persons with no temporomandibular (jaw) pain. The proportion with sleep disturbance among persons with jaw pain increased with the severity of the pain, rising from 20 percent for those with mild pain, to 32 percent for moderate pain, and 59 percent for severe pain.

Psychosocial Dimensions

The social and psychological dimensions of well-being and quality of life are deeply intertwined in everyday life and so are considered together here. Findings are reported for those oral conditions where the most research has been conducted: dental diseases (primarily those that affect appearance or involve extensive tooth loss), pain conditions, craniofacial deformities (primarily clefting syndromes), and oral and pharyngeal cancers. Psychological distress associated with oral health problems has been measured through individual questions (e.g., items assessing worry) and through standardized psychological instruments such as the Beck Depression Inventory (Beck 1967) or the Symptom Checklist 90 (SCL-90) (Derogatis 1983). Overall, the results point to a poorer quality of life and a tendency to avoid social contact as a result of concerns over facial appearance. Persistent pain has similar isolating and depressing effects.

Cultural Significance of Teeth

Cultural beliefs and customs are a major influence on the psychosocial effects of dental disease on individuals. In a historical essay on the art of pulling teeth, Kunzle (1989) noted that "the tooth has always been accorded a special, even magical, role among all peoples and at all times, and has stood for power, and its loss for loss of power." The practice of hiding or burying a lost tooth was based on the fear that its

recovery by an enemy could be used magically to inflict harm on the tooth owner. Remnants of that belief persist in the ritual of the tooth fairy, who rescues the hidden baby tooth and leaves a reward. The potency of teeth is well recorded in the metaphors of language. As Ziolkowski (1976) observed, "if something sets our teeth on edge or if someone casts an insult in our teeth, we can gnash our teeth in anger, show our teeth belligerently, grit our teeth resolutely, take the bit in our teeth, arm ourselves to the teeth, and fight tooth and nail in the teeth of great danger and, with luck, escape by the skin of our teeth."

The perception of healthy teeth contrasts sharply with that of diseased teeth. The sixteenth-century French surgeon Ambroise Paré remarked that "toothache was the greatest and most eternal of all pains . . . the fiery torture of the damned in hell" (Kunzle 1989); and a preacher of that time saw the root of original sin in a rotting tooth, as quoted in Kunzle (1989): "We unfortunate humans: We all have toothache and suffer ever and always from the teeth with which Adam bit the forbidden apple." In the nineteenth century, Sir James Frazer reported that African tribal kings could not be crowned if they were symbolically emasculated by having a broken tooth, a theme Freud echoed in declaring that dreams of pulling teeth were symbols of castration (Ziolkowski 1976).

The link between diseased teeth and weakness, impotence, and even moral turpitude and sin has been analyzed by scholars exploring the vast legacy of dental themes in art and literature. These authors offer a variety of political, social, psychological, and economic interpretations of diseased versus healthy teeth, attesting to the seriousness with which humanity has invested dental pain and tooth loss.

Cultures have sanctioned a variety of alterations to teeth by shaping and filing, embedding with jewels, bleaching, capping, or providing orthodontic treatments to improve occlusion and aesthetics. These procedures have been variously designed to enhance the status, power, and attractiveness of the owner. Cultures have also dictated specific practices to indicate social position, such as the former Japanese custom of dyeing a married woman's teeth black to denote her marital status.

Decisions about aesthetic surgery (Kaw 1993), the definition of what constitutes a severe malocclusion, and the need for dental aesthetics (Cons et al. 1986) and surgical intervention (Strauss 1985) depend on social norms. Recently, researchers have begun to examine the perceived attractiveness of the human smile in terms of tooth length, shape, and color, lip line and plumpness, and tooth exposure

(Dunn et al. 1996). Although there are no reliable studies to document the proportion of people who would seek treatment for perceived defects, the growth of practitioner groups and dental supply companies that specialize in aesthetic dentistry attests to a substantial increase in cosmetic dental treatments. Follow-up studies of patients who have undergone tooth bleaching indicate high levels of satisfaction with the extent and stability of the color change (Haywood et al. 1994).

Dental Problems and Social Function

Given the importance of the mouth and teeth in verbal and nonverbal communication, diseases that disrupt their functions are likely to damage self-image and alter the ability to sustain and build social relationships. The social functions of individuals encompass a variety of roles, from the most intimate in dating and mating behaviors, to other interpersonal contacts, to participation in social or community activities. Dental diseases and disorders can interfere with these social roles at any or all levels (Patrick and Bergner 1990). Whether because of social embarrassment or functional problems, people with oral conditions may avoid conversation or laughing, smiling, or other nonverbal expressions that show their mouth and teeth.

Two surveys of different segments of the adult population in Ontario, Canada, addressed the issue of self-consciousness or embarrassment related to oral health problems. Both inquired into the frequency of such problems in the prior year. Among persons over 18 years old in North York, Ontario, 7 percent reported limiting conversation with others because of oral problems in the prior year, 15 percent reported that they had avoided laughing or smiling because of oral problems, and 19 percent reported being embarrassed at least sometimes by the appearance or health of their teeth or mouth (Locker and Miller 1994). Among persons 50 years of age and older throughout the province, 24 percent reported being self-conscious about their appearance at least occasionally because of problems with their teeth, mouth, or dentures, 22 percent reported feeling uncomfortable at least occasionally, and 13 percent reported that they avoided smiling (Locker and Slade 1993).

Recently, investigators have begun to develop quantitative tools to assess patterns of facial animation, including normal facial expressions, in order to assess functional impairment in these movements (Trotman et al. 1998)—and ultimately to address implications for quality of life. The effects of facial

expressions, particularly smiling, on social relationships warrant further research.

The more subtle aspects of social function may also be affected by relatively common oral conditions such as tooth loss. In a study of elderly people with tooth loss in the United Kingdom, 30 percent reported difficulty chewing. Although only 5 percent had changed their diet, 9 percent felt uncomfortable chewing in front of others, and 13 percent reported embarrassment during social interactions (Smith and Sheiham 1979). Bergendal (1989) acknowledged tooth loss in adults to be a serious life event with a more difficult period of readjustment than retirement. In a study by Fiske et al. (1998), 25 percent of people without teeth reported that they had avoided close relationships because of fear of rejection when their toothlessness was discovered. The authors reported that reactions to tooth loss by the elderly include lowered self-confidence and altered self-image, bereavement, dislike of appearance, and the perception of being more advanced in age. A national study surveying 3,000 homes in Great Britain regarding the effect of oral health on the quality of life was performed in 1997-98. Seventy percent of respondents reported that their oral health affected their quality of life—either positively or negatively. Older people in higher socioeconomic groups, especially those who had seen a dentist within the last year, were more likely to report an enhancing effect (McGrath and Bedi 1998). In a study of older Californians, people who viewed themselves as more attractive than other people their age were more likely to have a greater number of natural teeth. They were also more likely to report less emotional anxiety, to rate their health as excellent or very good, and to be less likely to wear full or partial removable dentures (Matthias et al. 1993).

Less severe oral conditions may also have adverse effects on social function. Table 6.2 summarizes self-reported levels of impact on personal contact and social integration from surveys of numerous population groups within the United States. Generally, the prevalence of adverse effects ranges from 1 to 10 percent, although 20 percent of senior center residents in California reported feeling uncomfortable when eating with others. Avoidance of smiling or conversation tended to be more frequent than limitation of social interaction. However, the largest amount of variation appeared to be among population subgroups, even within studies. For example, within the Boston area, Veterans Affairs ambulatory care patients in the Veterans Health Study had substantially higher prevalence rates of

adverse effects than community-dwelling, healthy men enrolled in the Normative Aging Study.

The effects on people's perceived ability to perform their usual social roles appear to be more subtle (Table 6.2). Depending on the phrasing of questions and the population subgroup surveyed, the prevalence of limitation in social roles varied from 0 to 5 percent, although as many as 10 percent of Veterans Affairs patients reported that their daily activities were affected because of oral conditions.

There are very few studies of the effects of dental treatment on specific forms of social function. One study of handicapped adults in the United Kingdom did, however, report large improvements in five aspects of social functioning following provision of

general dental care, including prosthodontic care (Fiske et al. 1990).

Impacts of Oral-Facial Pain

Acute pain from dental caries can usually be treated effectively or, as in the case of recurrent aphthous ulcers (canker sores), will resolve over time. Chronic pain in the oral-facial region presents a different situation in that the cause may not be removable and treatment focuses on pain management. Researchers have compared chronic oral-facial pain to other types of pain in terms of severity and emotional impact. In a study of adult enrollees of a large health maintenance organization, Von Korff et al. (1988)

TABLE 6.2
Reported impacts of oral conditions on social function among U.S. population groups

Social Function Dimension(s) Assessed	Percentage of Population Group Reporting Effect		
	Good Effect	Bad Effect	No Effect
Intimacy			
Elderly Persons in North Carolina ^a			
Perceived effect of mouth on:			
Sex appeal	75	21	5
Kissing	78	18	4
Romantic relationships	81	17	3
Personal contact/social integration			
Persons Aged 18 to 61 in Five U.S. Sites ^b			
Avoid conversation	4		
Personal contact/social integration			
Elderly Persons in California ^c			
Reported impacts due to oral conditions:			
Senior Center Residents		Medicare Enrollees	
Limiting contact with people	10	1	
Feeling uncomfortable eating with others	20	4	
Personal contact/social integration			
Elderly Persons in Florida ^d			
Avoid laughing or smiling	6		
Avoid conversation	3		
Social roles			
Elderly Persons in North Carolina ^e			
Limit activities			
5			
Selected Populations in Boston Area^e			
Normative Aging Study Veterans Health Study Women's Health Project			
Personal contact/social integration			
Avoidance of conversation	2	10	7
Social interaction affected	1	9	7
Social roles			
Daily activities affected	1	10	10
Elderly Persons in North Carolina^f			
African Americans Whites			
Personal contact/social integration			
Avoided smiling	10	3	
Unable to enjoy people's company	5	1	
Avoided going out with others	3	0	
Less tolerant of others	3	0	
Social roles			
Difficulty doing jobs	1	0	
Unable to work	3	0	

^a Strauss and Hunt 1993.

^d Gilbert et al. 1993.

^b Gooch et al. 1989.

^e Kressin 1997.

^c Atchison and Dolan 1990.

^f Hunt et al. 1995.

compared pain in the temporomandibular region to back pain, headache, chest pain, and abdominal pain on a number of dimensions. The usual intensity of temporomandibular disorder (TMD) pain (4.3 on a 10-point scale) was similar to that of chest pain (4.3) and back pain (4.7); abdominal pain and headache were rated as more intense (5.1 and 5.9, respectively). TMD pain and back pain were more persistent than other pains, with 28 percent of persons with TMD pain (and 29 percent of those with back pain) reporting pain on more than half of the days in the prior 6 months. The daily duration of pain was also relatively high for TMD pain, with 27 percent reporting pain for 9 or more hours per day; this proportion was roughly the same as for headache and back pain. In this setting, persons with TMD pain sought care for their pain at about the same rate as persons with other pain, with about one quarter having sought care in the past 6 months and about 60 percent having sought care at some time in the past.

Locker and Grushka (1987) reported that more than 70 percent of persons with all types of acute and chronic pain worried about their dental health. Using scales of the Symptom Checklist 90 (SCL-90), age- and sex-standardized to the population under study, persons in the general population who reported TMD pain had substantially higher levels of anxiety and depression than those without a current pain condition (Von Korff et al. 1988). In the same study, items from the SCL-90 were used to assign an algorithm for diagnosis of depression. The percentage of persons with TMD who met the criteria for a possible diagnosis of major depression was 11 percent, compared with 2 percent among those without a current pain condition and 3 to 5 percent in the general population. Recent studies suggest that depression in TMD patients follows the onset of symptoms and reflects uncertainty about the cause(s) and the lack of effective treatments.

Social Responses to Facial Appearance

Just as cultural considerations color an individual's response to dental disease and tooth loss, so, too, does culture play a major role in the psychosocial impact of craniofacial deformities. More than any other body part, the face bears the stamp of individual identity, a fact neurobiology confirms in identifying an area of the brain that is dedicated to the recognition of faces.

In ancient Greece the face was seen as the mortal reflection of the gods, and those faces deemed most beautiful were also judged to be morally superior. Although cultures differ in the details, there appear

to be some invariant factors in the judgment of facial beauty and deformity (Stafford et al. 1989), and they are learned early in life. One U.S. study concluded that perceptions of what does and does not constitute an attractive face are established by age 7 (Cross and Cross 1971).

Aristotle also thought that the face, because of its position in the uppermost part of the body, pointed to the highest, most spiritual parts of the cosmos. Something of this mystical belief is preserved in linguistic traditions that regard what is "higher" and "ascending" as nobler than what is "lower" and "descending," whether in reference to body parts, human feelings, or classes and castes in society.

But if facial beauty is the mark of truth, moral superiority, nobility, and the soul, what of the opposite? Faces judged ugly or disharmonious, or those marred or scarred by birth defects or injury, have been associated with defects in character, intelligence, and morals. There is a long tradition of reading character on the basis of facial and head shapes according to one or another school of "physiognomics." Some schools likened human faces to animal species, such as cats, monkeys, or horses, endowing the human with the traits of the animal in question. Others constructed facial geometries that purported to show which races were nobler, which closer to animal ancestors. During the nineteenth century, Cesare Lombroso, an Italian physician and criminologist, developed a measuring scheme linking certain facial types to criminality (Magli 1989).

Given such a tradition, it is not surprising that people continue to "judge a book by its cover." A large body of research indicates that attractiveness has an important effect on psychological development and social relationships (Berscheid 1980). Noteworthy among early social psychological studies was the demonstration that physical attractiveness plays a key role in social expectations (Clifford and Walster 1973). Schoolteachers who were asked to evaluate the educational potential of students from school records and facial photographs judged attractive students to have higher educational potential and social ability than unattractive students. This finding has been replicated in a variety of school settings and grade levels.

The process of attributing positive characteristics to physically attractive persons is called the "beauty is good hypothesis." The positive impact of attractive appearance has been demonstrated to influence a wide range of social activities, from legal proceedings (Sigall and Ostrove 1975) to psychotherapeutic prognosis. This body of research indicates that appearance affects social expectations not only in educa-

tional performance, but also in vocational, marital, legal, and health care endeavors. However, the concept may not apply to all personal attributes. Bennett and Stanton (1993) found that appearance may have little effect on perceptions of intelligence or honesty.

Discrimination and stigmatization have historically characterized social responses to deformities. People whose appearance is significantly different from what society considers "normal" continue to be stigmatized. Examples in literature abound, in fable and myth, novels, movies, and plays (e.g., *Beauty and the Beast*, *The Elephant Man*, *The Hunchback of Notre Dame*, *The Phantom of the Opera*, and *Scarface*). Goffman (1963) provided a theory of stigmatization useful in understanding social responses to human differences and health conditions (Ablon 1981). Persons seen as deviant experience social devaluation associated with prejudices about the causes and consequences of congenital and acquired deformities (Macgregor 1974).

Ethnographic studies of facial deformities demonstrate marked variation in how birth defects are explained and treated (Cheng 1990, Meyerson 1990, Schepher-Hughes 1990, Toliver-Weddington 1990). Many cultures have regarded birth defects and the appearance of multiple births as ill omens, with the infants fated to be abandoned or killed. Even today, when it is understood that birth defects may result from an inherited genetic disorder, decisions about what that means and what if anything should be done about it are affected by religious or spiritual beliefs (Strauss 1988).

Progress in dentistry, medicine, and surgery has raised expectations that deformed or injured persons can be treated to enhance their appearance. Although critics have objected to the high value that U.S. society places on attractiveness and youthful appearance, and on how this emphasis influences opportunity, advocates of remediation counter that social values are hard to change and that individuals must adapt to a society's norms and stereotypes. Corrective surgery does produce positive assessments from others (Berscheid and Gangestad 1982). People in postoperative photographs following reconstructive and plastic surgery were seen as kinder, more sexually appealing, more likely to be better marriage partners, and more employable and successful than the same individuals in preoperative views.

Cleft Lip/Palate and Malocclusion

Persons with cleft lip/palate have a variety of problems related to appearance, eating, and speech. Studies of self-concept (Broder and Strauss 1989, Kapp-Simon 1986), psychosocial development (Richman and Eliason 1982, Richman et al. 1988), and social perception by peers (Schneiderman and Harding 1984, Tobiasen 1987), parents and teachers (Mitchell et al. 1984, Schneiderman and Auer 1984), and the public (Middleton et al. 1986) highlight the psychosocial problems of children with cleft lip/palate (Broder and Richman 1987).

These problems often continue into adolescence, with appearance and speech remaining problematic (Richman et al. 1985), even when the individual has had comprehensive care (Strauss et al. 1988). Studies show an association between cleft lip/palate and the increased reporting of conduct problems at home (Tobiasen and Hiebert 1984) and of behavioral and learning problems at school (Tobiasen et al. 1987). Although children with cleft lip/palate have not been shown to suffer from a negative self-concept (Kapp 1979, Richman 1983) or psychological disorders, they often become socially inhibited and self-conscious (Richman and Eliason 1982).

Adolescent girls with cleft lip/palate have higher rates of social adjustment problems, particularly related to appearance (Leonard et al. 1991). Using psychological projection methods, one study (Pillemer and Cook 1989) also concluded that children with facial deformities may have inhibited personality styles as well as reduced expectations for success in social interactions. Kapp-Simon (1986) found that social relationships negatively influence overall self-esteem and that adjustment of adolescents is associated with the degree of inhibition (Kapp-Simon et al. 1992).

Many of the earliest studies of the consequences of oral disease examined the impact of developmental disorders such as facial clefts and malocclusion on personal contact and social integration. There are few outcome studies that demonstrate how social, marital, and occupational status may be affected in adults with facial deformities or cleft lip/palate. In studies dating from the 1970s, persons with cleft lip/palate achieved greater educational levels than did their fathers, but did not exceed their fathers' occupational status (McWilliams and Paradise 1973). Persons with cleft lip/palate were less upwardly mobile than their siblings and achieved lower adult incomes (Peter et al. 1975). Although persons with cleft

lip/palate had higher occupational desires, they had lower income expectations than did controls; adults felt socially inept and had a tendency to shy away from group activities (Van Demark and Van Demark 1970). Subtle forms of discrimination and stigmatization were experienced among adults with cleft lip/palate (McWilliams 1970).

Studies of marital status have indicated that persons with cleft lip/palate are less likely than nonaffected peers to date (Peter et al. 1975) or marry (McWilliams and Paradise 1973), and that when they marry, they marry later (Heller et al. 1981, Peter and Chinsky 1974). Persons with cleft lip/palate were also more likely to remain childless or have fewer children (Peter and Chinsky 1974).

The evidence concerning anatomical misalignments of the jaws (e.g., overbite, open bite, crossbite) is less consistent, with some early studies showing no clear relationship between the malocclusion and job opportunities or social discrimination (Shaw et al. 1980a). However, people with severe malocclusions are likely to be teased and to have difficulty interacting socially. These problems improve following orthodontic and orthognathic treatment (Cunningham et al. 1996, Helm et al. 1985, Shaw et al. 1980b).

Oral and Pharyngeal Cancers

Surgical treatment for oral and pharyngeal cancers can result in functional impairment as well as permanent disfigurement. Problems may include the loss of part of the tongue, loss of taste, loss of chewing ability, difficulty in speaking, and pain. Furthermore, in addition to concerns about their function and their future, oral and pharyngeal cancer patients must cope with an altered appearance. In a study of patients who were disease-free from 6 months to 8 years following surgical tumor removal, Gamba et al. (1992) reported that those with more pronounced disfigurement had greater changes in self-image, a worsened relationship with their partner, reduced sexuality, and increased social isolation. A study by Bjordal and Kaasa (1995) also noted that 30 percent of oral and pharyngeal cancer patients were still experiencing psychological distress 7 to 11 years after treatment. Depression, too, is frequent in cancer patients. Patients with oral and pharyngeal cancers are at an even greater risk for depression than other cancer patients, due to surgeries that alter their appearance (Gritz et al. 1999). Because oral and pharyngeal cancers are also frequently associated with chronic alcohol and tobacco use, depression may be related to withdrawal from these substances or to

preexisting psychopathology. Persistent pain, as noted earlier, may also be a contributing factor to depression.

Gritz et al. (1999) conducted a prospective analysis of changes in quality of life over time with the aim of identifying which factors might be predictive of future improvements or declines. Participants were 186 oral and pharyngeal cancer patients, all smokers or recent former smokers, diagnosed with primary carcinomas of the oral cavity, pharynx, or larynx. The patients were tested at baseline, at 1 month after radiation and/or surgery, and 1 year later (for a subset of 105 patients available for follow-up). Measures used included the Karnofsky Performance Scale, which uses expert judgments of functional performance scored from 0 to 100; the Cancer Rehabilitation Evaluation System Short Form, in which patients rate their quality of life along physical, psychosocial, marital, sexual, and medical interaction scales; the previously mentioned Performance Status Scale for Head and Neck Cancer Patients (which includes scales for eating and speaking); and the Profile of Mood States, in which patients rate their feelings over the previous week, yielding analyses that enable scaling along six mood states: tension-anxiety, depression-dejection, anger-hostility, confusion-bewilderment, and vigor-activity. Results indicated that in spite of functional improvement on some scales over time, there was continued dysfunction in speech and eating. Patients also reported declines in marital and sexual functioning, as well as an increase in alcohol use. Interestingly, the best predictor of quality of life 1 year after treatment was the scores obtained after initial smoking cessation advice was given, while the patients were undergoing treatment and in recovery. Other predictors were treatment type (quality of life was generally poorer for radiation patients) and score on the vigor subscale of the Profile of Mood States. The investigators concluded that medical follow-up must integrate tailored psychological and behavioral interventions to achieve better quality of life for oral and pharyngeal cancer patients.

Indirect Economic Costs

The financial impact of oral disease on quality of life is easiest to quantify in terms of direct per capita costs of oral disease and treatment as well as the costs of publicly supported dental care programs. The Health Care Financing Administration includes these costs in its annual total health care expenditure reports. These expenditures are presented in Chapter 9, and the related data on utilization of care are given

in Chapter 4. Estimating the indirect costs associated with oral health conditions and treatments, such as disability days or lost productivity, is more difficult. Several methodologies are used to estimate these costs, although relatively few studies have actually estimated the indirect costs associated with mortality or lost productivity due to dental conditions or treatments (Reisine and Locker 1995).

The annual National Health Interview Survey routinely collects data on disability days associated with selected conditions. As shown in Table 6.3, there were 3.7 days of restricted activity per 100 employed persons 18 years and older reported in the United States in 1996 associated with an acute dental condition, as defined by a dental symptom or treatment visit. Restricted activity days were most prevalent among adults aged 18 to 24 years, women, blacks, and individuals with annual incomes of less than \$10,000. Compared to the 624.0 restricted activity days per 100 persons per year for all acute conditions, the 3.7 restricted activity days for dental conditions represent a relatively small loss on an individual basis. They do, however, add up to a sizable number of days lost from work or school every year for the population as a whole.

In addition, Table 6.3 presents the number of bed days and work-loss days per 100 employed persons aged 18 and older. Also, for youths 5 to 17 years of age, 3.1 days of school were lost per year.

An important aspect of identifying the specific health-related quality of life burden of oral disease will be to acquire more data on how the overall health-related quality of life of persons with oral disease differs from that of healthy persons, those with other diseases and activity limitations, and those with co-morbid oral and general health problems. The CDC's "Healthy Days" measures and population

data—used by all states since 1993 on the Behavioral Risk Factor Surveillance System (BRFSS) and recently added to the National Health and Nutrition Examination Survey (NHANES)—offer the potential of examining quality of life outcomes in relation to oral health and other disease measures used in these surveys (Gift 1996). These productivity-related measures ask about the number of recent days when physical health was not good, mental health was not good, and activities were limited. These measures were found to be valid in use with general populations and among older persons and adults with disabilities (Moriarty and Zack 1999). Related measures on activity limitations and recent days of pain, depression, anxiety, sleeplessness, and vitality are also asked by about half of all BRFSS states and—when used with oral disease measures—could help to identify the impact of oral disease prevention programs at the state and local levels.

In the 1996 National Health Interview Survey, the percentage of all acute conditions that are medically attended for all ages is 67.9 percent. "Medically attended" is defined as having contacted a physician (or other provider) or having a condition that causes a person to cut back on activities for at least half a day. For acute dental conditions, 59.6 percent are medically attended for all ages (NCHS 1996).

The most dramatic oral diseases by virtue of their high mortality rates are oral and pharyngeal cancers. The Centers for Disease Control and Prevention estimated in 1988 that 16.2 years of life were lost per person dying of cancer of the oral cavity and pharynx (CDC 1991). This exceeds the average for all cancer sites, which was 15.4 years lost.

Researchers are beginning to assess costs associated with chronic craniofacial conditions such as periodontal diseases, pain syndromes, and congenital anomalies. These estimates may also include the costs to caregivers who take time off from work to attend to these needs or to take children to the dentist. Recent estimates put the lifetime costs of multiple surgeries and the other medical, dental, and rehabilitation therapies typical of the team approach to the habilitation of individuals with cleft lip or palate at a minimum of \$101,000 (Waitzman et al. 1996). Overall costs of chronic pain conditions in America are estimated to be \$79 billion (Bonica 1990). Given the prevalence of headaches and

TABLE 6.3
Disability days due to all acute conditions and acute dental conditions,
United States, 1996

	All Acute Conditions		Acute Dental Conditions	
	Total Days (in thousands)	Days per 100 Persons	Total Days (in thousands)	Days per 100 Persons
School-loss days ^a	152,305	296.9	1,611	3.1 ^b
Work-loss days ^c	358,377	284.0	2,442	1.9 ^b
Bed days ^d	717,868	271.7	4,602	1.7 ^b
Restricted activity ^d	1,648,932	624.0	9,705	3.7 ^b

^a Youths 5 to 17 years of age.

^b Figure does not meet NCHS standard of reliability or precision.

^c Currently employed persons 18 years and older.

^d Persons of all ages.

Source: NCHS 1996.

temporomandibular disorders, the amount representing chronic oral-facial pain would certainly be in the billions.

RATINGS OF ORAL HEALTH

Researchers use two ways to assess how individuals rate their oral health: global ratings and satisfaction ratings. Global ratings employ a ranking scale, with excellent health at one end and poor health at the other. Satisfaction ratings are more akin to oral-health-related quality of life measures insofar as they ask individuals how satisfied (or dissatisfied) they are with their oral health status in relation to symptoms, physical function, appearance, social function, and psychological status.

Global Ratings

The dental survey conducted in connection with the International Collaborative Survey II (ICS II) asked younger (35 to 44 years) and older adults (65 to 74 years) of various ethnic groups at three sites in the United States to rate their oral health from 1 (excellent) to 6 (poor). The vast majority of younger and older white adults rated their oral health as good to excellent (82 and 80 percent, respectively, in Baltimore, and 71 and 68 percent, respectively, in San Antonio). In contrast, American Indians, Hispanics, and African Americans were more likely to rate their oral health as fair or poor. More detailed analysis found a significant relationship between oral health ratings and perceived general health, dentate status, importance of oral health, income, oral pain, symptoms, and dental visits (Chen et al. 1997).

Studies in the United States (Bloom et al. 1992, Chen et al. 1997, Gift et al. 1997, Matthias et al. 1995) have related global ratings of oral health to demographic, clinical, and psychosocial factors. A special supplement on oral health in the National Health Interview Survey in 1989 asked participants to rate their oral health as excellent, very good, good, fair, or poor (Bloom et al. 1992). The majority of people (67 percent) rated their oral health as excellent or very good, 23 percent rated it good, and only 10 percent rated it fair or poor. Men, younger people, and more frequent users of dental services tended to rate their oral health better than women, older adults, and less frequent users. African Americans, Hispanics, and American Indians were less likely to rate their oral health positively than were whites in similar geographic locations (Atchison and Gift 1997).

A study of Medicare participants in California (Matthias et al. 1995) also found that when asked to

rate oral health as excellent, very good, good, fair, or poor, most people rated it as good or above (74 percent). Interestingly, the relationship between self-rated global oral health and clinical measures used to rate the severity of dental caries or periodontal disease was weak. The finding of such inconsistency has been confirmed in other studies (Gooch et al. 1989, Reisine and Bailit 1980, Rosenberg et al. 1988) and was noted by Wilson and Cleary (1995). The most important correlates of self-rated oral health were worry about teeth and appearance, race, education, general health status, and depression scores.

Satisfaction Ratings

According to Jokovic and Locker (1997), "expressions of satisfaction and dissatisfaction are important oral health status indicators since they synthesize objective health states, subjective responses, and culturally based values and expectations." ICS I represented an early effort at assessing satisfaction with dental status (Arnlijot and WHO 1985). Specifically, it looked at satisfaction with teeth and gingiva among adults and adolescents in metropolitan and non-metropolitan areas in 10 countries. With the exception of Japanese participants, satisfaction with teeth and gingiva among adolescents was fairly high, but the percentage of adults satisfied with teeth among participants from all countries was relatively low. A decade later, ICS II (Chen et al. 1997) again assessed whether younger and older adults were satisfied with the way their teeth looked. American Indians were the least satisfied with the appearance of their teeth. Other community surveys of satisfaction (Barenthin 1977, Gilbert et al. 1994, Jokovic and Locker 1997, Murtomaa and Laine 1985, Rosenoer and Sheiham 1995, Van Waas et al. 1994) show that most people are satisfied with the performance and appearance of their teeth.

ORAL-HEALTH-RELATED QUALITY OF LIFE MEASURES

Much of the research on oral health and quality of life focuses on the negative impact of craniofacial diseases and disorders, diverting attention from the positive effects of good oral health. Although it could be argued that the benefits are self-evident, few studies have investigated how people value oral health. Strauss and Hunt (1993) found that older adults in North Carolina felt that the presence of teeth enhanced their appearance, ability to eat, and enjoyment of food, and that teeth also had a positive effect on comfort, confidence, speech, enjoyment, and

longevity. More generally, a large multisite study in the United States found that most adults across ethnic groups believe in the seriousness of oral disease and the importance of oral health for general health (Davidson et al. 1996, Nakazono et al. 1997).

During the last decade, researchers have identified and described specific effects of oral disease on quality of life, and there are now at least 11 questionnaires designed to measure oral-health-related quality of life (Slade et al. 1998). These range from the three-item Rand Dental Health Index (pain, worry, conversation) to the 49-item Oral Health Impact Profile, which includes items relating to function, pain, physical disability, psychological disability, social disability, and handicap.

Other researchers have used generic quality of life questionnaires (Reisine and Weber 1989) or analyzed oral health survey data, statistically summarizing clinical measures with self-reported symptoms, perceptions, and behaviors to create scales of oral-health-related quality of life (Gift et al. 1997, Gilbert et al. 1997). An essential requirement for these analyses is the establishment of theoretical models that describe and distinguish among quality of life dimensions.

Although the questionnaires have been used in population studies, there is increased emphasis on the need to incorporate concepts of quality of life into outcome research, using the questionnaires in longitudinal or intervention studies to examine changes in quality of life following provision of dental care. Individual items from the questionnaires can be used in describing the impact of oral health status on specific functions. For example, the questions concerning eating dysfunction presented in Table 6.1 are derived from such questionnaires. Multidimensional indicators also can be used to produce quantitative scores that indicate the severity of the impact caused by oral health problems. Because there are seldom any reference values or population norms to indicate thresholds at which such scores represent "abnormal" or "severe" impacts on quality of life, the results are more valuable for analytic studies examining trends and associations than for describing impacts of oral health within populations.

Surveys in the United States and elsewhere have used multidimensional questionnaires and have revealed consistent correlations between reduced quality of life and poorer clinical oral status and reduced access to dental care. Table 6.4 provides an overview of these studies, their populations, and descriptions of the assessment tools used. In the studies using varied tools, there are consistent relationships between quality of life and standard epi-

demiological indices of missing teeth, decayed teeth, and periodontal disease—conditions that are linked to a lack of dental care. In several studies, infrequent dental visits and problem-motivated dental visits were found to be independently associated with reduced quality of life in multivariate models (Gilbert et al. 1997, Slade and Spencer 1994a). Lower socioeconomic status was an additional explanatory factor in five studies (Atchison and Dolan 1990, Gilbert et al. 1997, Gooch et al. 1989, Leao and Sheiham 1995, Locker and Slade 1994).

All of the studies in Table 6.4 used cross-sectional designs to analyze associations with quality of life; it is thus not possible to infer a causal link between quality of life and the factors under study. Information about change in quality of life from those questionnaires is available from only two population-based longitudinal studies. In one, a study of elderly Floridians, small and statistically nonsignificant changes in the Geriatric Oral Health Assessment Index were observed following an oral health promotion program (Dolan 1997). In the second, a 2-year longitudinal study of elderly South Australians, change in quality of life was measured using the Oral Health Impact Profile. The study did not find that regular dental attendees had consistently better patterns of change than episodic dental attendees (Slade and Spencer 1994b).

In contrast, some information from experimental studies of carefully selected clinical samples indicates effects of specific dental treatments on quality of life. Using the Sickness Impact Profile in a U.S. study of 30 patients treated for TMDs, Reisine et al. (1989) found significant improvements for rest/sleep, home tasks, work, and leisure following treatment. A randomized clinical trial of 63 Canadians treated for TMDs compared changes in quality of life, measured using visual analog scales. The investigators found equivalent improvements in quality of life, which mirrored improvements in pain ratings, in all three treatment groups (Dao et al. 1994). A German study by Schliephake et al. (1996) reported improvements in quality of life, measured using a 22-item scale, among 85 patients following tumor removal and reconstructive surgery.

Two multidimensional scales have been specifically designed to assess global disability (e.g., disability days, pain interference with daily activities) related to recurrent or chronic pain. Both the Graded Chronic Pain Scale (Von Korff et al. 1992) and the Multidimensional Pain Inventory (Kerns et al. 1985) have been used to assess disability related to TMDs, as well as back pain and headache. Although TMD patients are generally found to have lower mean levels

of disability than patients with other types of pain using these scales, both scales can discriminate the full range of disability in TMDs. For example, the Graded Chronic Pain Scale is an ordinal scale that assigns persons with pain to hierarchical grades indicating low-intensity, nondisabling pain (grade I); high-intensity, nondisabling pain (grade II); high-disability, moderately limiting pain (grade III); and high-disability, severely limiting pain (grade IV). In a sample of HMO patients with TMDs, 41 percent were grade I, 43 percent grade II, 11 percent grade III, and 5 percent grade IV (Von Korff et al. 1992). That is, about 16 percent of patients seeking care for TMDs were moderately or severely limited by their pain.

HEIGHTENED EXPECTATIONS

Advances in public health and biomedical research in the late twentieth century have transformed our lives remarkably, adding years to the life span, providing cures for many diseases and disorders, and promising a new era of genetic medicine and bioengineering. The new field of science aimed at formulating con-

cepts and methods for determining health-related quality of life is a reflection of these advances. People living in societies that enjoy the benefits of this progress are responding with heightened expectations: They want to live long and live well, free of infirmity, impairment, disability, and handicap; they want an optimal quality of life and well-being. In terms of oral health, new social norms and cultural values now dictate that teeth should be retained over the lifetime, and oral pain and dysfunction forever banished. Quality of life measures applied to oral health outcomes will be used to further those goals.

FINDINGS

Examination of efforts to characterize the functional and social implications of oral and craniofacial diseases reveals the following findings:

- Oral health is related to well-being and quality of life as measured along functional, psychosocial, and economic dimensions. Diet, nutrition, sleep, psychological status, social interaction, school, and work are affected by impaired oral and craniofacial health.

TABLE 6.4
Studies of multidimensional quality of life measures

	Population Studied	Numbers of Individuals	Assessment Tool	Description of Assessment Tool
Gooch et al. 1989	U.S. insured adults aged 18-61 years	902 female 756 male	Rand Dental Health Index	Three dental questions written to represent factors contributing to adverse effects of dental disease
Atchison and Dolan 1990	California Medicare recipients aged 65+	1,000 female 755 male	Geriatric Oral Health Assessment Index (GOHAI)	A series of 12 questions measuring patient-reported oral functional problems
Hunt et al. 1995	N.C. elderly aged 70+	440	Oral Health Impact Profile (OHIP)	A comprehensive measure of self-reported dysfunction consisting of 49 questions
Kressin et al. 1996	Male veterans aged 47+	1,242 male	Oral Health-related Quality Of Life (OHQOL)	A brief global assessment of the impact of oral conditions consisting of three items
Gift et al. 1997	U.S. aged 18+	760 female 555 male	1981 Health Resources and Services Administration study	Multidimensional concept using data from a large national sample
Gilbert et al. 1997	Floridians aged 45+	491 female 383 male	Oral Disadvantage Assessment	Eight self-reported measures of avoidance in daily activities due to decrements in oral health
Locker and Miller 1994	Canadians aged 18+	299 female 244 male	Subjective Oral Health Status Indicators	Five oral health status indicators based on WHO's International Classification of Impairments, Disabilities and Handicaps
Locker and Slade 1994	Canadians aged 50+	168 female 144 male	Oral Health Impact Profile (OHIP)	A comprehensive measure of self-reported dysfunction consisting of 49 questions
Slade and Spencer 1994b	Australians aged 60+	660 female 557 male	Oral Health Impact Profile (OHIP)	A comprehensive measure of self-reported dysfunction consisting of 49 questions
Leao and Sheiham 1995	Brazilians aged 35-44	303 female 359 male	Dental Impact on Daily Living (DIDL)	Thirty-six questions that assess the oral health impacts on daily living
Coates et al. 1996	Australian dental patients	635 (+795 previously surveyed)	Oral Health Impact Profile (OHIP)	A comprehensive measure of self-reported dysfunction consisting of 49 questions

- Cultural values influence oral and craniofacial health and well-being and can play an important role in care utilization practices and in perpetuating acceptable oral health and facial norms.

- Oral and craniofacial diseases and their treatment place a burden on society in the form of lost days and years of productive work. Acute dental conditions contribute to a range of problems for employed adults, including restricted activity, bed days, and work loss, and school loss for children. In addition, conditions such as oral and pharyngeal cancers contribute to premature death and can be measured by years of life lost.

- Oral and craniofacial diseases and conditions contribute to compromised ability to bite, chew, and swallow foods; limitations in food selection; and poor nutrition. These conditions include tooth loss, diminished salivary functions, oral-facial pain conditions such as temporomandibular disorders, alterations in taste, and functional limitations of prosthetic replacements.

- Oral-facial pain, as a symptom of untreated dental and oral problems and as a condition in and of itself, is a major source of diminished quality of life. It is associated with sleep deprivation, depression, and multiple adverse psychosocial outcomes.

- Self-reported impacts of oral conditions on social function include limitations in verbal and non-verbal communication, social interaction, and intimacy. Individuals with facial disfigurements due to craniofacial diseases and conditions and their treatments may experience loss of self-image and self-esteem, anxiety, depression, and social stigma; these in turn may limit educational, career, and marital opportunities and affect other social relations.

- Reduced oral-health-related quality of life is associated with poor clinical status and reduced access to care.

REFERENCES

- Ablon J. Stigmatized health conditions. *Soc Sci Med [Med Anthropol]* 1981 Jan;15B(1):5-9.
- Aday LA, Forthofer RN. A profile of black and Hispanic subgroups' access to dental care: findings from the National Health Interview Survey. *J Public Health Dent* 1992;52(4):210-5.
- Agerberg G, Carlsson GE. Symptoms of functional disturbances of the masticatory system. A comparison of frequencies in a population sample and in a group of patients. *Acta Odontol Scand* 1975;33(4):183-90.
- Andersen RM, Davidson PL. Ethnicity, aging and oral health outcomes: a conceptual framework. *Adv Dent Res* 1997 May;11(2):203-9.
- Arnlijot HA, World Health Organization (WHO), editors. *Oral health care systems: an international collaborative study*. London: Quintessence; 1985.
- Atchison KA, Dolan TA. Development of the Geriatric Oral Health Assessment Index. *J Dent Educ* 1990 Nov;54(11):680-7.
- Atchison KA, Gift HC. Perceived oral health in a diverse sample. *Adv Dent Res* 1997 May;11(2):272-80.
- Barenthin I. Dental health status and dental satisfaction. *Int J Epidemiol* 1977 Mar;6(1):73-9.
- Beck AT. *Depression: clinical, experimental, and theoretical aspects*. New York: Hoeber; 1967.
- Bennett ME, Stanton ML. Psychotherapy for persons with craniofacial deformities: can we treat without theory? *Cleft Palate Craniofac J* 1993 Jul;30(4):406-10.
- Bergendal B. The relative importance of tooth loss and denture wearing in Swedish adults. *Community Dent Health* 1989;6:103-11.
- Bergner M, Bobbitt RA, Carter WB, Gilson BS. The Sickness Impact Profile: development and final revision of a health status measure. *Med Care* 1981 Aug;19(8):787-805.
- Berscheid E. Overview of the psychological effects of physical attractiveness. In: Lucker GW, Ribbens KA, McNamara JA, editors. *Psychological aspects of facial form*. Proceedings of a sponsored symposium honoring Professor Robert E. Moyers; 1980 Feb and Mar; Ann Arbor: Center for Human Growth and Development, The University of Michigan; 1980, 1981. Craniofacial growth series; monograph no. 11. p. 1-23.
- Berscheid E, Gangestad S. The social psychological implications of facial physical attractiveness. *Clin Plast Surg* 1982 Jul;9(3):289-96.
- Bjordan K, Kaasa S. Psychological distress in head and neck cancer patients 7-11 years after curative treatment. *Br J Cancer* 1995 Mar;71(3):592-7.
- Blaum CS, Fries BE, Fiatorone MA. Factors associated with low body mass index and weight loss in nursing home residents. *J Gerontol Med Sci* 1995;50:M162-8.
- Bloom B, Gift HC, Jack SS. Dental services and oral health. *Vital Health Stat* 10 1992 Dec;(183):1-95.
- Bonica JJ. General considerations of chronic pain. In: Bonica JJ, Loeser JD, Chapman CR, Fordyce WE, editors. *The management of pain*. Vol. 1. Philadelphia: Lea and Febiger; 1990. p. 180-96.
- Broder HL, Richman LC. An examination of mental health services offered by cleft/craniofacial teams. *Cleft Palate J* 1987 Apr;24(2):158-62.
- Broder HL, Strauss RP. Self-concept of early primary school age children with visible or invisible defects. *Cleft Palate J* 1989 Apr;26(2):114-7.
- Brodeur JM, Laurin D, Vallee R, Lachapelle D. Nutrient intake and gastrointestinal disorders related to masticatory performance in the edentulous elderly. *J Prosthet Dent* 1993 Nov;70(5):468-73.

- Carlsson GE. Masticatory efficiency: the effect of age, the loss of teeth and prosthetic rehabilitation. *Int Dent J* 1984 Jun;34(2):93-7.
- Centers for Disease Control (CDC). Cigarette smoking among adults—United States, 1988. *JAMA* 1991 Dec 11;266(22):3113-4.
- Chauncey HH, Muench ME, Kapur KK, Wayler AH. The effect of the loss of teeth on diet and nutrition. *Int Dent J* 1984;34(2):98-104.
- Chen M, Andersen RM, Barmes DE, Leclercq MH, Lyttle CS. Comparing oral health care systems—a second international collaborative study. Geneva: World Health Organization; 1997.
- Cheng LR. Asian-American cultural perspectives on birth defects: focus on cleft palate. *Cleft Palate J* 1990 Jul;27(3):294-300.
- Clifford M, Walster E. The effect of physical attractiveness on teacher expectations. *Soc Educ* 1973;46:248-58.
- Coates E, Slade GD, Goss AN, Gorkic E. Oral conditions and their social impact among HIV dental patients. *Aust Dent J* 1996 Feb;41(1):33-6.
- Cons NC, Jenny J, Kohout FJ. DAI: the dental aesthetic index. Iowa City: University of Iowa College of Dentistry; 1986.
- Cross JF, Cross J. Age, sex, race and the perception of facial beauty. *Dev Psychol* 1971;5:433-9.
- Cunningham SJ, Crean SJ, Hunt NP, Harris M. Preparation, perceptions, and problems: a long-term follow-up study of orthognathic surgery. *Int J Adult Orthodon Ortognath Surg* 1996;11(1):41-7.
- Dao TT, Lavigne GJ, Charbonneau A, Feine JS, Lund JP. The efficacy of oral splints in the treatment of myofascial pain of the jaw muscles: a controlled clinical trial. *Pain* 1994 Jan;56(1):85-94.
- Davidson PL, Andersen RM. Determinants of dental care utilization for diverse ethnic and age groups. *Adv Dent Res* 1997 May;11(2):254-62.
- Davidson PL, Andersen RM, Marcus M, Atchison KA, Reifel N, Nakazono T, Rana H. Indicators of oral health in diverse ethnic and age groups: findings from the International Collaborative Study of Oral Health Outcomes (ICS-II) USA research locations. *J Med Syst* 1996 Oct;20(5):295-316.
- Demers M, Bourdages J, Brodeur JM, Benigeri M. Indicators of masticatory performance among elderly complete denture wearers. *J Prosthet Dent* 1996 Feb;75(2):188-93.
- Derogatis LR. Symptom Checklist-90. Administration, scoring and procedures manual for the revised version. Baltimore: Clinical Psychometric Research; 1983.
- Diehnelt D, Kiyak HA, Beach BH. Predictors of oral health behaviors among elderly Japanese-Americans. *Spec Care Dentist* 1990;10:112-8.
- Dolan T. The sensitivity of the Geriatric Oral Health Assessment Index to dental care. *J Dent Educ* 1997;61(1):37-46.
- Dunn WJ, Murchison DF, Broome JC. Esthetics: patients' perceptions of dental attractiveness. *J Prosthodont* 1996 Sep;5(3):166-71.
- Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, Kessler RC. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA* 1998 Nov 11;280(18):1569-75.
- Ernest SL. Dietary intake, food preferences, simulated salivary flow rate, and masticatory ability in older adults with complete dentitions. *Spec Care Dentist* 1993 May-Jun;13(3):102-6.
- Febres C, Echeverri EA, Keene HJ. Parental awareness, habits, and social factors and their relationship to baby bottle tooth decay. *Pediatr Dent* 1997 Jan-Feb;19(1):22-7.
- Feldman RS, Kapur KK, Alman JE, Chauncey HH. Aging and mastication: changes in performance and in the swallowing threshold with natural dentition. *J Am Geriatr Soc* 1980 Mar;28(3):97-103.
- Fiske J, Gelbier S, Watson RM. The benefit of dental care to an elderly population assessed using a socio-dental measure of oral handicap. *Br Dent J* 1990;168(4):153-6.
- Fiske J, Davis DM, Frances C, Gelbier S. The emotional effects of tooth loss in edentulous people. *Brit Dent J* 1998;184(2):90-3.
- Fries JF, Spitz PW, Young DY. The dimensions of health outcomes: the health assessment questionnaire, disability and pain scales. *J Rheumatol* 1982 Sep-Oct;9(5):789-93.
- Gamba A, Romano M, Grosso IM, Tamburini M, Cantu G, Molinari R, Ventafridda V. Psychosocial adjustment of patients surgically treated for head and neck cancer. *Head Neck* 1992 May Jun;14(3):218-23.
- Garrett NR, Kapur KK, Perez P. Effects of improvements of poorly fitting dentures and new dentures on patient satisfaction. *J Prosthet Dent* 1996 Oct;76(4):403-13.
- Geertman ME, Boerrigter EM, Van't Hof MA, Van Waas MA, van Oort RP, Boering G, Kalk W. Two-center clinical trial of implant-retained mandibular overdentures versus complete dentures—chewing ability. *Community Dent Oral Epidemiol* 1996 Feb;24(1):79-84.
- Gift HC. Quality of life—an outcome of oral health care? *Public Health Dent* 1996;56(2):67-8.
- Gift HC, Atchison KA. Oral health, health, and health-related quality of life. *Med Care* 1995 Nov;33(11 Suppl):NS57-77.
- Gift HC, Atchison KA, Dayton CM. Conceptualizing oral health and oral health-related quality of life. *Soc Sci Med* 1997 Mar;44(5):601-8.
- Gilbert GH, Heft MW, Duncan RP. Oral signs, symptoms, and behaviors in older Floridians. *J Public Health Dent* 1993 Summer;53(3):151-7.
- Gilbert GH, Heft MW, Duncan RP, Ringelberg ML. Perceived need for dental care in dentate older adults. *Int Dent J* 1994 Apr;44(2):145-52.

- Gilbert GH, Duncan RP, Heft MW, Dolan TA, Vogel WB. Oral disadvantage among dentate adults. *Community Dent Oral Epidemiol* 1997 Aug;25(4):301-13.
- Godson JH, Williams SA. Oral health and health related behaviours among three-year-old children born to first and second generation Pakistani mothers in Bradford, UK. *Community Dent Health* 1996 Mar;13(1):27-33.
- Goffman E. *Stigma: notes on the management of spoiled identity*. Englewood Cliffs (NJ): Prentice-Hall; 1963. p. 2-3.
- Gooch BF, Dolan TA, Bourque LB. Correlates of self-reported dental health status upon enrollment in the Rand Health Insurance Experiment. *J Dent Educ* 1989 Nov;53(11):629-37.
- Goulet JP, Lavigne GJ, Lund JP. Jaw pain prevalence among French-speaking Canadians and related symptoms of temporomandibular disorders. *J Dent Res* 1995 Nov;74(11):1738-44.
- Gritz ER, Carmack CL, de Moor C, Coscarelli A, Schacherer CW, Meyers EG, Abemayor E. First year after head and neck cancer: quality of life. *J Clin Oncol* 1999;17(1):352-60.
- Guyatt AH, Feeny DH, Patrick DL. Measuring health-related quality of life. *Ann Intern Med* 1993 Apr 15;118(8):622-9.
- Harris EF, Woods MA, Robinson QC. Dental health patterns in an urban Midsouth population: race, sex and age changes. *Quintessence Int* 1993 Jan;24(1):45-52.
- Haywood VB, Leonard RH, Nelson CF, Brunson WD. Effectiveness, side effects and long-term status of nightguard vital bleaching. *J Am Dent Assoc* 1994 Sep;125(9):1219-26.
- Helkimo E, Carlsson GE, Helkimo M. Chewing efficiency and state of dentition. A methodologic study. *Acta Odontol Scand* 1978;36(1):33-41.
- Heller A, Tidmarsh W, Pless IB. The psychosocial functioning of young adults born with cleft lip or palate. A follow-up study. *Clin Pediatr* 1981;20:459-65.
- Helm S, Kreiborg S, Solow B. Psychosocial implications of malocclusion: a 15-year follow-up study in 30-year-old Danes. *Am J Orthod* 1985 Feb;85(2):110-8.
- Henderson GE, et al., editors. *The social medicine reader*. Durham (NC): Duke University Press; 1997. p. 6-11.
- Hildebrandt GH, Dominguez BL, Schork MA, Loesche WJ. Functional units, chewing, swallowing, and food avoidance among the elderly. *J Prosthet Dent* 1997 Jun;77(6):588-95.
- Hufford DJ. Folk medicine in contemporary America. In: Kirkland J, et al., editors. *Herbal and magical medicine*. Durham: Duke University Press; 1992. p. 14-31.
- Hunt RJ, Slade GD, Strauss RP. Differences between racial groups in the impact of oral disorders among older adults in North Carolina. *J Public Health Dent* 1995 Fall;55(4):205-9.
- Hunt SM, McEwen J, McKenna SP. Measuring health status: a new tool for clinicians and epidemiologists. *J R Coll Gen Pract* 1985 Apr;35(273):185-8.
- Idowu AT, Graser GN, Handelman SL. The effect of age and dentition status on masticatory function in older adults. *Spec Care Dentist* 1986 Mar-Apr;6(2):80-3.
- Jokovic A, Locker D. Dissatisfaction with oral health status in an older adult population. *J Public Health Dent* 1997 Winter;57(1):40-7.
- Kapp K. Self-concept of the cleft lip and/or palate child. *Cleft Palate J* 1979 Apr;16(2):171-6.
- Kapp-Simon K. Self-concept of primary-school-age children with cleft lip, cleft palate, or both. *Cleft Palate J* 1986 Jan;23(1):24-7.
- Kapp-Simon KA, Simon DJ, Kristovich S. Self-perception, social skills, adjustment, and inhibition in young adolescents with craniofacial anomalies. *Cleft Palate Craniofac J* 1992 Jul;29(4):352-6.
- Kaw E. Medicalization of racial features: Asian-American women and cosmetic surgery. *Med Anthropol Q* 1993;7(1):74-89.
- Kayser AF. Shortened dental arches and oral function. *J Oral Rehabil* 1981 Sep;8(5):457-62.
- Keller HH. Malnutrition in institutionalized elderly: how and why? *J Am Geriatr Soc* 1993 Nov;41(11):1212-8.
- Kelly M, Bruerd B. The prevalence of baby bottle tooth decay among two Native American populations. *J Public Health Dent* 1987 Spring;47(2):94-7.
- Kerns RD, Turk DC, Rudy TE. The West Haven-Yale Multidimensional Pain Inventory (WHYMPI). *Pain* 1985 Dec;23(4):345-56.
- Kiyak HA. Age and culture: influences on oral health behavior. *Int Dent J* 1993 Feb;43(1):9-16.
- Kleinman A. *Sickness as cultural semantics: issues for an anthropological medicine and psychiatry*. In: Ahmed P, Coelho G, editors. *Toward a new definition of health: psychosocial dimensions*. New York: Plenum; 1979. p. 53-65.
- Kolasa K. I won't cook turnip greens if you won't cook kielbasa: food behavior or Polonia and its health implications. In: Bauwens E, editor. *The anthropology of health*. St. Louis: Mosby; 1978. p. 130.
- Kressin NR. Oral health-related quality of life measure (OHQOL). In: Slade GD, editor. *Measuring oral health and quality of life*. Chapel Hill: University of North Carolina, School of Dental Ecology; 1997.
- Kressin N, Spiro A 3rd, Bosse R, Garcia R, Kazis L. Assessing oral health-related quality of life: findings from the normative aging study. *Med Care* 1996 May;34(5):416-27.
- Kunzle D. The art of pulling teeth in the seventeenth and nineteenth centuries: from public martyrdom to private nightmare and political struggle? Part 3. In: Feher M, with Nadaff R, Razi N, editors. *Cambridge (MA): distributed by MIT Press; 1989. p. 28-89.*
- Leake JL. An index of chewing ability. *J Public Health Dent* 1990 Summer;50(4):262-7.

- Leao A, Sheiham A. Relation between clinical dental status and subjective impacts on daily living. *J Dent Res* 1995 Jul;74(7):1408-13.
- Lee J, Kiyak HA. Oral disease beliefs, behaviors, and health status of Korean-Americans. *J Public Health Dent* 1992 Spring;52(3):131-6.
- Lee KL, Schwarz E, Mak KY. Improving oral health through understanding the meaning of health and disease in a Chinese culture. *Int Dent J* 1993 Feb;43(1):2-8.
- Leonard BJ, Brust JD, Abrahams G, Sielaff B. Self-concept of children and adolescents with cleft lip and/or palate. *Cleft Palate Craniofac J* 1991 Oct;28(4):347-53.
- List MA, Ritter-Sterr C, Lansky SB. A performance status scale for head and neck cancer patients. *Cancer* 1990;66:564-9.
- Locker D, Grushka M. The impact of dental and facial pain. *J Dent Res* 1987 Sep;66(9):1414-7.
- Locker D, Miller Y. Subjectively reported oral health status in an adult population. *Community Dent Oral Epidemiol* 1994 Dec;22(6):425-30.
- Locker D, Slade G. Prevalence of symptoms associated with temporomandibular disorders in a Canadian population. *Community Dent Oral Epidemiol* 1988 Oct;16(5):310-3.
- Locker D, Slade G. Oral health and the quality of life among older adults: the Oral Health Impact Profile. *J Can Dent Assoc* 1993 Oct;59(10):830-3,837-8,844.
- Locker D, Slade G. Association between clinical and subjective indicators of oral health status in an older adult population. *Gerodontology* 1994 Dec;11(2):108-14.
- Macgregor FC. Transformation and identity, the face and plastic surgery. New York: Quadrangle/New York Times; 1974. p. 119.
- Magli P. The face and the soul. In: Feher M, with Nadaff R, Razi N, editors. *Fragments for a history of the human body. Part 2.* Cambridge (MA): distributed by MIT Press; 1989. p. 87-127.
- Marcus SE, Drury TF, Brown LJ, Zion GR. Tooth retention and tooth loss in the permanent dentition of adults: United States, 1988-1991. *J Dent Res* 1996 Feb;75(Spec No):684-95.
- Matthias RE, Atchison KA, Schweitzer SO, Lubben JE, Mayer-Oakes A, De Jong F. Comparisons between dentist ratings and self-ratings of dental appearance in an elderly population. *Spec Care Dentist* 1993 Mar-Apr;13(2):53-60.
- Matthias RE, Atchison KA, Lubben JE, De Jong F, Schweitzer SO. Factors affecting self-ratings of oral health. *J Public Health Dent* 1995;55(4):197-204.
- McEwen B. Stress and the brain. In: Conlan R, editor. *States of mind.* New York: John Wiley; 1999. p. 81-102.
- McGrath C, Bedi R. A study of the impact of oral health on the quality of life of older people in the UK—findings from a national survey. *Gerodontology* 1998;15(2):93-8.
- McWilliams BJ. Psychosocial development and modification. Washington: American Speech and Hearing Association; 1970. ASHA Reports No. 5.
- McWilliams BJ, Paradise LP. Educational, occupational, and marital status of cleft palate adults. *Cleft Palate J* 1973 Jul;10:223-9.
- Meenan RF, Gertman PM, Mason JH. Measuring health status in arthritis. The arthritis impact measurement scales. *Arthritis Rheum* 1980 Feb;23(2):146-52.
- Meyerson MD. Cultural considerations in the treatment of Latinos with craniofacial malformations. *Cleft Palate J* 1990 Jul;27(3):279-88.
- Middleton GF, Lass NJ, Starr P, Pannbacker M. Survey of public awareness and knowledge of cleft palate. *Cleft Palate J* 1986 Jan;23(1):58-62.
- Mitchell CK, Lott R, Pannbacker M. Perceptions about cleft palate held by school personnel: suggestions for in-service training development. *Cleft Palate J* 1984 Oct;21(4):308-12.
- Moriarty D, Zack M. Validation of the Centers for Disease Control and Prevention's Healthy Days measures [abstract]. *Qual Life Res* 1999;8(7):617.
- Murtomaa H, Laine P. Differentiating dental satisfaction in Finns by means of discriminant analyses. *Community Dent Oral Epidemiol* 1985 Feb;13(1):7-10.
- Nakazono TT, Davidson PL, Andersen RM. Oral health beliefs in diverse populations. *Adv Dent Res* 1997 May;11(2):235-44.
- National Center for Health Statistics (NCHS). Current estimates from the National Health Interview Survey, 1996. Series 10, no. 200. Hyattsville (MD): Public Health Service; 1996.
- Navia JM. Carbohydrates and dental health. *Am J Clin Nutr* 1994 Mar;59(Suppl 3):719S-275.
- Patrick DL, Bergner M. Measurement of oral health status in the 1990s. *Annu Rev Public Health* 1990;11:165-83.
- Peter JP, Chinsky RR. Sociological aspects of cleft palate adults: I. Marriage. *Cleft Palate J* 1974 Jul;11(0):295-309.
- Peter JP, Chinsky RR, Fisher MJ. Sociological aspects of cleft palate adults: III. Vocational and economic aspects. *Cleft Palate J* 1975 Apr;12:193-9.
- Pillemer FG, Cook KV. The psychosocial adjustment of pediatric craniofacial patients after surgery. *Cleft Palate J* 1989 Jul;26(3):201-7, Discussion 207-8.
- Ranta K, Tuominen R, Paunio I. Perceived oral health status and ability to chew among an adult Finnish population. *Gerodontology* 1987 Jun;3(3):136-9.
- Reisine ST, Bailit HL. Clinical oral health status and adult perceptions of oral health. *Soc Sci Med [Med Psychol Med Sociol]* 1980 Dec;14A(6):597-605.
- Reisine S, Locker D. Social, psychological and economic impact of oral conditions and treatments. In: Cohen LK, Gift HC, editors. *Disease prevention and oral health promotion: socio-dental sciences in action.* Copenhagen: Munksgaard; 1995. p. 33-72.